

Central-to-peripheral nuclear modification factors in Pb–Pb collisions at $\sqrt{s_{\text{NN}}} = 17.3$ GeV

NA57 Collaboration

F. Antinori^k, P.A. Bacon^e, A. Badalà^f, R. Barbera^f, A. Belogianni^a, I.J. Bloodworth^e, M. Bombara^h, G.E. Bruno^{b,*}, S.A. Bull^e, R. Caliendo^b, M. Campbell^g, W. Carena^g, N. Carrer^g, R.F. Clarke^e, A. Dainese^{k,*}, D. Di Bari^b, S. Di Libertoⁿ, R. Divià^g, D. Elia^b, D. Evans^e, G.A. Feofilov^p, R.A. Fini^b, P. Ganoti^a, B. Ghidini^b, G. Grella^o, H. Helstrup^d, K.F. Hetland^d, A.K. Holme^j, A. Jacholkowski^f, G.T. Jones^e, P. Jovanovic^e, A. Jusko^e, R. Kamermans^r, J.B. Kinson^e, K. Knudson^g, V. Kondratiev^p, I. Králik^h, A. Kravčákováⁱ, P. Kuijer^r, V. Lenti^b, R. Lietava^e, G. Løvhøiden^j, V. Manzari^b, M.A. Mazzoneⁿ, F. Meddiⁿ, A. Michalon^q, M. Morando^k, P.I. Norman^e, A. Palmeri^f, G.S. Pappalardo^f, B. Pastirčák^h, R.J. Platt^e, E. Quercigh^k, F. Riggi^f, D. Röhrich^c, G. Romano^o, K. Šafařík^g, L. Šándor^h, E. Schillings^r, G. Segato^k, M. Sené^l, R. Sené^l, W. Snoeys^g, F. Soramel^{k,l}, M. Spyropoulou-Stassinaki^a, P. Staroba^m, R. Turrisi^k, T.S. Tveter^j, J. Urbánⁱ, P. van de Ven^r, P. Vande Vyvre^g, A. Vascotto^g, T. Vik^j, O. Villalobos Baillie^e, L. Vinogradov^p, T. Virgili^o, M.F. Votruba^e, J. Vrlákováⁱ, P. Závada^m

^a Physics Department, University of Athens, Athens, Greece

^b Dipartimento IA di Fisica dell'Università del Politecnico and INFN, Bari, Italy

^c Fysisk Institutt, Universitetet i Bergen, Bergen, Norway

^d Høgskolen i Bergen, Bergen, Norway

^e School of Physics and Astronomy, University of Birmingham, Birmingham, UK

^f University of Catania and INFN, Catania, Italy

^g CERN, European Laboratory for Particle Physics, Geneva, Switzerland

^h Institute of Experimental Physics, Slovak Academy of Science, Košice, Slovakia

ⁱ P.J. Šafařík University, Košice, Slovakia

^j Fysisk Institutt, Universitetet i Oslo, Oslo, Norway

^k University of Padua and INFN, Padua, Italy

^l Collège de France, Paris, France

^m Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic

ⁿ University "La Sapienza" and INFN, Rome, Italy

^o Dipartimento di Scienze Fisiche "E.R. Caianiello" dell'Università and INFN, Salerno, Italy

^P State University of St. Petersburg, St. Petersburg, Russia

^Q Institut de Recherches Subatomique, IN2P3/ULP, Strasbourg, France

[†] Utrecht University and NIKHEF, Utrecht, The Netherlands

Received 2 May 2005; accepted 15 July 2005

Available online 1 August 2005

Editor: M. Doser

Abstract

We present central-to-peripheral nuclear modification factors, R_{CP} , for the p_T distributions of K_S^0 , Λ , $\bar{\Lambda}$, and negatively charged particles, measured at central rapidity in Pb–Pb collisions at top SPS energy. The data cover the 55% most central fraction of the inelastic cross section. The K_S^0 and Λ $R_{CP}(p_T)$ are similar in shape to those measured at $\sqrt{s_{NN}} = 200$ GeV at RHIC, though they are larger in absolute value. We have compared our K_S^0 R_{CP} data to a theoretical calculation. The prediction overestimates the data at $p_T \approx 3\text{--}4$ GeV/ c , unless sizeable parton energy loss is included in the calculation.

© 2005 Elsevier B.V. All rights reserved.

PACS: 12.38.Mh; 25.75.Nq; 25.75.Dw

Keywords: Nucleus–nucleus collisions; Strange particles

1. Introduction

The quenching of high transverse momentum (p_T) particles in central heavy-ion collisions is one of the main discoveries at BNL-RHIC. The effect is quantified using the nuclear modification factor:

$$R_{AA}(p_T) = \frac{1}{\langle N_{\text{coll}} \rangle_C} \times \frac{d^2 N_{AA}^C / d p_T dy}{d^2 N_{pp} / d p_T dy}, \quad (1)$$

where $\langle N_{\text{coll}} \rangle_C$ is the average number of nucleon–nucleon (NN) collisions for nucleus–nucleus (AA) collisions in a given centrality class C. The nuclear modification factor would be equal to unity if the AA collision were a mere superposition of N_{coll} independent nucleon–nucleon collisions. In central Au–Au collisions at a c.m.s. energy per nucleon–nucleon pair of $\sqrt{s_{NN}} = 200$ GeV, the PHENIX and STAR experiments have measured a suppression by a factor 4–5 with respect to unity in R_{AA} for $p_T \gtrsim 5$ GeV/ c , independent of the particle species [1]. A similar suppression at high p_T is observed also in the central-to-

peripheral nuclear modification factor

$$R_{CP}(p_T) = \frac{\langle N_{\text{coll}} \rangle_P}{\langle N_{\text{coll}} \rangle_C} \times \frac{d^2 N_{AA}^C / d p_T dy}{d^2 N_{AA}^P / d p_T dy}, \quad (2)$$

where a class P of peripheral nucleus–nucleus collisions replaces the pp reference (see, e.g., [2]). The measured suppression is interpreted as being due to energy loss of the hard partons traversing the high-density QCD medium expected to be formed in high-energy heavy-ion collisions [3]. Parton energy loss would predominantly occur via the mechanism of medium-induced gluon radiation [4,5].

Preliminary results [6] from the RHIC run at $\sqrt{s_{NN}} = 62.4$ GeV show a R_{AA} suppression by about a factor 3, not much smaller than that observed at $\sqrt{s_{NN}} = 200$ GeV. This motivates the search for parton energy loss effects at even lower c.m.s. energy, i.e., in Pb–Pb collisions with $\sqrt{s_{NN}} = 17.3$ GeV at CERN-SPS. Possible indications of such effects were found by the WA98 Collaboration [7], who observed a suppression of the π^0 R_{CP} . In addition, a recent π^0 R_{AA} analysis [8], using the WA98 Pb–Pb data [7] and a parameterization of a wide compilation of pp data at similar energy available in the literature, favours a scenario of significant energy loss.

We investigate the presence of energy loss effects in Pb–Pb collisions at $\sqrt{s_{NN}} = 17.3$ GeV by mea-

* Corresponding authors. Correspondence to G.E. Bruno, via Orabona 4, I-70126 Bari, Italy, correspondence to A. Dainese, via Marzolo 8, I-35131 Padova, Italy.

E-mail addresses: giuseppe.bruno@ba.infn.it (G.E. Bruno), andrea.dainese@pd.infn.it (A. Dainese).

¹ Permanent address: University of Udine, Udine, Italy.

suring $R_{CP}(p_T)$ for K_S^0 , Λ and $\bar{\Lambda}$ particles, and for unidentified negatively charged particles, h^- . The relative behaviour of the $R_{CP}(p_T)$ patterns for K_S^0 and Λ particles is also expected to be sensitive to parton coalescence effects [9] in the hadronization dynamics. It has been suggested that these effects occur at RHIC energy (see, e.g., [10]).

In Section 2 the experimental setup, the data sets and the analysis procedures are described. The R_{CP} results are presented in Section 3 and they are compared in Section 4 to other experimental results at SPS and RHIC energies as well as with theoretical calculations.

2. Apparatus, data sets and analysis

The NA57 apparatus, described in detail in [11], was designed to study the production of strange and multi-strange particles in fixed-target heavy-ion collisions by reconstructing their weak decays into final states containing charged particles only. Tracks are reconstructed in the 30 cm-long silicon telescope: an array of pixel detector planes with a cross section of $5 \times 5 \text{ cm}^2$ placed inside a 1.4 T magnetic field normal to the beam direction. The telescope is inclined in the non-bending plane by a 40 mrad angle relative to the beam line and it points to the target located 60 cm from the first detector plane. The acceptance covers about half a unit in rapidity at central rapidity and transverse momentum larger than about 0.5 GeV/c. For the Pb–Pb runs, the centrality trigger, based on charged multiplicity, was set so as to select approximately the most central 60% of the inelastic collisions.

The results presented in this Letter are based on the analysis of data collected during the 1998 (about 2.1×10^8 events) and 2000 (about 2.5×10^8 events) Pb–Pb runs with beam momentum of 158 A GeV/c.

The collision centrality is determined using the charged particle multiplicity N_{ch} in the pseudorapidity range $2 < \eta < 4$, sampled by the microstrip silicon detectors (MSD) as described in [12,13]. N_{ch} is related to the centrality assuming $N_{ch} = q \times N_{part}^\alpha$ (a modified wounded nucleon model) [12], where N_{part} is the number of participants, i.e., nucleons participating in the primary nucleon–nucleon collisions, estimated from the Glauber model [14]. The N_{ch} distribution is well described by a wounded nucleon model fit with $\alpha =$

Table 1

Average number of participants and of NN collisions with their systematic errors, as a function of centrality

Class (% σ_{inel}^{Pb-Pb})	$\langle N_{part} \rangle$	$\langle N_{coll} \rangle$
0.0–5.0%	345.3 ± 1.7	779.2 ± 26.6
10.0–20.0%	214.7 ± 5.8	421.7 ± 26.1
20.0–30.0%	143.0 ± 6.6	247.7 ± 21.5
30.0–40.0%	92.6 ± 6.4	140.5 ± 16.2
40.0–55.0%	49.5 ± 5.0	63.8 ± 9.8

1.0 [13]. The inelastic cross-section extracted from the fit [12], using an inelastic non-diffractive NN cross section $\sigma_{inel}^{NN} = 30 \text{ mb}$ [15,16], is $\sigma_{inel}^{Pb-Pb} = 7.26 \text{ b}$. Five centrality classes are defined, with N_{ch} limits corresponding to given fractions of σ_{inel}^{Pb-Pb} . For each class the average values of N_{part} and N_{coll} are calculated from the Glauber model, with Pb nucleus Woods–Saxon density-profile parameters as given in [17]. In Table 1 we present the definitions of the five centrality classes along with the corresponding values of $\langle N_{part} \rangle$ and $\langle N_{coll} \rangle$ with their systematic errors, estimated by varying the α parameter in the range [1.0, 1.1], the Woods–Saxon parameters within their tabulated uncertainties and the inelastic non-diffractive NN cross section at $\sqrt{s} \simeq 17 \text{ GeV}$ within its systematic error estimated to be 1.5% [15,16].² Note, however, that the contribution due to the uncertainty on σ_{inel}^{NN} is negligible for $\langle N_{part} \rangle$, while for $\langle N_{coll} \rangle$ it ranges from 1.5% in the most central class to 1.1% in the most peripheral, but it cancels out in the ratio $\langle N_{coll} \rangle_P / \langle N_{coll} \rangle_C$ used for R_{CP} .

Strange particles are reconstructed using their decay channels into charged particles: $K_S^0 \rightarrow \pi^+ \pi^-$, $\Lambda \rightarrow \pi^- p$, and $\bar{\Lambda} \rightarrow \pi^+ \bar{p}$. The selection procedure is described in detail in [18,19]. The main criteria are the following: (a) the two decay tracks are compatible with the hypothesis of having a common origin point; (b) the reconstructed secondary vertex is well separated in space from the target; (c) the reconstructed candidate points back to the primary vertex position. For K_S^0 , the two decay tracks are required to miss the interaction vertex by applying a p_T -dependent cut on the product of their impact parameters (distances

² The $\langle N_{part} \rangle$ values given in Table 1 for the most central and the most peripheral classes differ slightly from those published in [13] due to a refined treatment of the trigger effect in the fit of the multiplicity distribution done for the present analysis.




Fig. 1. Invariant mass distributions for K_S^0 (upper row) and Λ (lower row) candidates in different p_T ranges. Arrows indicate the nominal masses of the two particles and shaded areas indicate the ranges considered for the analysis. No centrality selection is applied.

of closest approach to the interaction vertex in the bending plane). Ambiguities among K_S^0 , Λ and $\bar{\Lambda}$ are eliminated by means of cuts [18] on the Podolanski–Armenteros plot [20]. The final invariant mass distributions for K_S^0 and Λ candidates in different p_T ranges are shown in Fig. 1. The shaded areas correspond to the windows used for this analysis; the size of these windows is smaller at low p_T , where the invariant mass resolution is better. The statistics amounts to 1.8×10^6 K_S^0 , 0.7×10^6 Λ , and 0.1×10^6 $\bar{\Lambda}$. We have estimated and subtracted the combinatorial background using the event-mixing technique on a subsample of events representative of the full statistics [19]. The background fraction in the selected mass window is negligible for our most peripheral class (40–55%) and it increases going to more central classes, where it also increases from low to high transverse momentum. In the worst case, in class 0–5%, the subtracted background fraction amounts to: $(6 \pm 2)\%$ for K_S^0 with $p_T > 2.5$ GeV/c; $(10 \pm 3)\%$ for Λ with $p_T > 2.5$ GeV/c; $(8 \pm 3)\%$ for $\bar{\Lambda}$ with $p_T > 2$ GeV/c.

Negatively charged particles, h^- , are selected from a sample of good-quality tracks (clusters in more than 80% of the telescope planes, less than 30% of the clusters shared with other tracks) using an impact parameter cut to ensure they come from the interaction vertex. The residual contamination of secondary tracks (decay products of weakly-decaying particles) has been estimated to be of about 2%, independent of p_T , for the most central class of events, and lower for the other classes. The statistics amounts to 10^8 particles.

We calculate $R_{CP}(p_T)$, Eq. (2), using p_T distributions which are unweighted, i.e., not corrected for geometrical acceptance and reconstruction/selection efficiency. For other analyses, in particular, for the measurement of the p_T -integrated strange particle production yields [21], we adopted a procedure, specifically developed for rarer signals like Ξ and Ω , in which every selected particle was assigned a correction weight, calculated on the basis of a Monte Carlo simulation [19]. This correction is very time consuming and, for the more abundant signals (h^- , K_S^0 and Λ), it was calculated only for a representative subsam-




Fig. 2. (Colour online.) Ratio of the average correction weights for a sample of central events (0–10%) and a sample of peripheral events (30–55%), as a function of p_T and of the particle species.

ple of the available statistics. For the present study, we have verified that these weights do not depend on the event centrality over the full transverse momentum range covered (see Fig. 2). Using unweighted spectra results in a negligible systematic error on $R_{CP}(p_T)$ as compared to the other contributions discussed in the following.

3. Results

We use class 40–55% as the reference peripheral class in the denominator of R_{CP} , see Eq. (2), and vary the ‘central’ class in the numerator from 0–5% to 30–40%. We estimated possible residual systematic effects by comparing the R_{CP} results corresponding to the year 1998 sample and to the year 2000 sample, which were also processed with different versions of the reconstruction software. We find compatible results, with point-to-point differences smaller than 10% for h^- , K_S^0 and $\Lambda + \bar{\Lambda}$. We assign ‘reconstruction’ systematic errors of 5% for all four species. For strange particles, a small contribution to the systematic errors has been recognized as due to the procedure used to remove fake tracks, which yield duplicated candidates, particularly in central collisions that have higher track density.³ For K_S^0 , Λ and $\bar{\Lambda}$, the resulting error, estimated to be at most 3% at high p_T for the ratio

³ For the h^- analysis, fake tracks were avoided by imposing tighter track-quality conditions.

0–5%/40–55% and smaller for the other ratios, has been added in quadrature to the ‘reconstruction’ systematic error and to the systematic error due to background subtraction, which is also at most 3%.

Fig. 3 shows our $R_{CP}(p_T)$ results for four different ‘central’ classes (C). The error bars are obtained as a quadratic sum of the statistical errors and the p_T -dependent systematic errors (< 7%). The shaded bands centered at $R_{CP} = 1$ represent the p_T -independent systematic error due to the uncertainty in the ratio $\langle N_{coll} \rangle_P / \langle N_{coll} \rangle_C$, while the shaded bands at low p_T represent the R_{CP} values corresponding to N_{part} -scaling, with the band indicating the systematic error due to the uncertainty in the ratio $\langle N_{part} \rangle_C / \langle N_{part} \rangle_P$. We first focus on the 0–5%/40–55% ratio. At low p_T (≈ 0.5 GeV/c), the h^- , K_S^0 and Λ patterns are compatible with N_{part} -scaling, while the $\bar{\Lambda}$ points are clearly below. As p_T increases, R_{CP} for the K_S^0 approaches one; R_{CP} for the Λ behaves differently from that of the K_S^0 above $p_T \approx 1$ GeV/c, reaching a value of about 1.5, as does the $\bar{\Lambda}$. For p_T values below 2 GeV/c, the R_{CP} for negative particles is dominated by negative pions, and stays below the corresponding values for K_S^0 . At higher p_T the h^- R_{CP} lies between those for K_S^0 and Λ ; in this region there may be significant K^- and \bar{p} contributions in the sample of negative particles (note that $\bar{p}/\pi^- \simeq 0.8$ for $p_T \gtrsim 3$ GeV/c in central Au–Au collisions at $\sqrt{s_{NN}} = 200$ GeV [22]). For less central collisions, the K_S^0 R_{CP} exhibit a small enhancement in the range $p_T \gtrsim 1.2$ GeV/c, while within errors the R_{CP} for the other particles do not vary with respect to 0–5%/40–55% R_{CP} .

The difference between the Λ and the $\bar{\Lambda}$ at low p_T , already observed in the centrality dependence of the p_T -integrated yields in Pb–Pb collisions [23], is pronounced. On the basis of the symmetry of the apparatus and of the signal extraction procedure, we are confident that the effect is not caused by an experimental bias. It may be due to a centrality-dependent absorption effect of $\bar{\Lambda}$ in a nucleon-rich environment. In this respect, note that at SPS energy the $\bar{\Lambda}/\Lambda$ ratio is measured to be lower by a factor about 2 in lead–lead with respect to sulfur-induced collisions [24] and the $\bar{\Lambda}$ yield per participant is measured to be lower by a factor 0.71 ± 0.05 in p–Pb [18] with respect to p–Be collisions [23].




Fig. 3. (Colour online.) Centrality dependence of $R_{CP}(p_T)$ for h^- , K_S^0 , Λ and $\bar{\Lambda}$ in Pb–Pb collisions at $\sqrt{s_{NN}} = 17.3$ GeV. Shaded bands centered at $R_{CP} = 1$ represent the systematic error due to the uncertainty in the ratio of the values of $\langle N_{coll} \rangle$ in each class; shaded bands at low p_T represent the values expected for scaling with the number of participants, together with their systematic error.

4. Comparisons and discussion

In Fig. 4 we compare our results to R_{CP} measurements at the SPS and at RHIC. In the left-hand panel, the WA98 π^0 data [7] for the ratio 1–6%/22–43% in Pb–Pb collisions at $\sqrt{s_{NN}} = 17.3$ GeV are plotted together with the NA57 h^- and K_S^0 data for the same centrality classes.⁴ Using these classes, the K_S^0 R_{CP} is approximately constant at 0.9 for $p_T > 1$ GeV/c and is significantly larger than that measured by the WA98 Collaboration for π^0 ($R_{CP} \approx 0.6$), even when taking into account the normalization systematic errors, independent for the two experiments. The h^- data from NA57 are compatible, within the system-

atic errors, with the π^0 data from WA98 for $p_T \lesssim 1.5$ GeV/c, where the h^- sample is expected to be dominated by π^- . For higher p_T , h^- have a larger R_{CP} than π^0 ; this may be due to increasing contributions from K^- and \bar{p} in the h^- sample. At top RHIC energy, $\sqrt{s_{NN}} = 200$ GeV, the kaon R_{CP} (as measured by PHENIX [25] and STAR [2]) is larger than that of neutral pions (PHENIX [25]) for $p_T \lesssim 2$ GeV/c, while they are similar for higher transverse momenta. The observed difference in R_{CP} between kaons and pions at SPS energy, and at RHIC energy for low p_T , is reminiscent of the ‘Cronin enhancement’ above N_{coll} -scaling originally observed for $3 < p_T < 6$ GeV/c in proton–nucleus (pA) collisions at $\sqrt{s_{NN}}$ values up to 38.8 GeV [26]. This enhancement, commonly interpreted as due to initial-state multiple scattering (partonic intrinsic transverse momentum broadening), was in fact found to increase according to the hierarchy pions–kaons–protons [26]. At RHIC energy, owing to the hardening of the p_T distributions, the effect would be reduced compared to lower energies, as confirmed

⁴ In the WA98 publication [7], the centrality classes are defined using percentiles of the measured minimum bias cross section, $\sigma_{WA98 m.b.}^{Pb-Pb} \approx 6.3$ b. Here, we rename the classes in terms of percentiles of the inelastic cross section, σ_{inel}^{Pb-Pb} . The values used by the two experiments for the inelastic cross section are very similar: 7.26 b for NA57 and 7.41 b for WA98.




Fig. 4. (Colour online.) Left: $R_{CP}(p_T)$ for h^- and K_S^0 from NA57 and π^0 from WA98 [7] in Pb–Pb collisions at $\sqrt{s_{NN}} = 17.3$ GeV. Right: $R_{CP}(p_T)$ for K_S^0 and Λ in Pb–Pb at $\sqrt{s_{NN}} = 17.3$ GeV (NA57) and in Au–Au at $\sqrt{s_{NN}} = 200$ GeV (STAR) [2]; slightly different peripheral classes are employed for the comparison. The bars centered at $R_{CP} = 1$ represent the normalization errors; the point-by-point bars are the quadratic sum of statistical and systematic errors.

by preliminary results on the particle-species dependence of the d–Au (similar to proton–nucleus) R_{CP} at $\sqrt{s_{NN}} = 200$ GeV [27].

The comparison for K_S^0 and Λ at SPS and RHIC (STAR data for Au–Au at $\sqrt{s_{NN}} = 200$ GeV [2]) is presented in the right-hand panel of Fig. 4. In the p_T range covered by our data, up to 4 GeV/c, the *relative* pattern for K_S^0 and Λ is similar at the two energies, while absolute values are higher at SPS than at RHIC, where parton energy loss is believed to have a strong effect. Part of the difference between the R_{CP} values at the two energies may be due to different nuclear modification of the parton distribution functions (PDFs). In fact, in the $x_{Bjorken}$ range relevant for parton production at a given p_T at the SPS, e.g., $x \simeq 0.3$ for $p_T \simeq 3$ GeV/c, nuclear PDFs are expected to be enhanced by about 10–20% (anti-shadowing), while almost no effect is expected for the smaller values, $x \simeq 0.03$, relevant for the same p_T values at RHIC energy (see, e.g., [28]). At RHIC, the larger R_{CP} for Λ with respect to kaons [2] or, more generally, for baryons with respect to mesons [25], in the intermediate p_T range, 2–4 GeV/c, has been interpreted as due to parton coalescence [9,10] in a high-density medium with partonic degrees of freedom. Our data show that a similar Λ –K pattern is present also at $\sqrt{s_{NN}} = 17.3$ GeV. We note that such a pattern may also be explained in terms of larger Cronin effect for Λ with respect to kaons.

In Fig. 5 we compare our K_S^0 data to predictions provided by X.N. Wang [29], obtained from a perturbative-QCD-based calculation [4,30], including (thick line) or excluding (thin line) in-medium parton energy loss. The initial gluonic rapidity density of the medium, dN_g/dy , was scaled down from that needed to describe RHIC data, according to the decrease by about a factor 2 in the charged multiplicity per unit of rapidity from RHIC to SPS energy. For the 0–5%/40–55% R_{CP} , the curve without energy loss shows a large enhancement, increasing with p_T . In the calculation this enhancement arises principally from initial-state partonic intrinsic transverse momentum broadening, which is assumed to be proportional to the number of scatterings that the two colliding partons suffer inside the nuclei before the hard scattering, and thus larger for central than for peripheral collisions [30]. The magnitude of the broadening is tuned on the basis of the original Cronin effect data [26]. A ‘Cronin-like’ enhancement is observed in the d–Au central-to-peripheral nuclear modification factor for charged kaons at RHIC energy [27], while it is not present in our nucleus–nucleus K_S^0 data. The curve that includes energy loss, scaled down from RHIC as explained above, describes the data better. Moving to less central collisions the predictions with and without energy loss get closer to one another, and both are compatible with the data. As a cross-check, we compare the R_{CP} value predicted by X.N. Wang




Fig. 5. (Colour online.) $R_{CP}(p_T)$ for K_S^0 in Pb–Pb collisions at $\sqrt{s_{NN}} = 17.3$ GeV, compared to predictions [29,31] with and without the effect of parton energy loss (details in the text).

with energy loss to the prediction of an independent model of parton energy loss, the parton quenching model (PQM), based on the BDMPS formalism [5], that describes several energy-loss-related observables at RHIC energies [31]. Since the PQM model does not include initial-state effects, the predicted R_{CP} was rescaled using the Wang baseline without energy loss (thin line). The PQM result has an uncertainty due to the fact that the medium-induced energy loss becomes of the order of the initial parton energy [31]; since this uncertainty is larger for low-energy partons, the calculation result is meaningful for $p_T \gtrsim 4$ GeV/c only. The PQM result, shown in Fig. 5 by the hatched areas, agrees with the Wang calculation with energy loss (thick line). The two models predict a similar energy loss effect at SPS energy, i.e., a reduction of the 0–5%/40–55% R_{CP} (for $p_T \gtrsim 4$ GeV/c) by about a factor 2, with respect to the value calculated without energy loss.

5. Conclusions

Central-to-peripheral nuclear modification factors for K_S^0 , Λ , $\bar{\Lambda}$ and h^- in Pb–Pb collisions at top SPS energy have been measured as a function of p_T up to about 4 GeV/c. At low p_T , R_{CP} agrees with N_{part} scaling for all the particles under consideration, except the $\bar{\Lambda}$, for which the yields at low p_T are found to increase slower than the number of participants. For $p_T \gtrsim 1$ GeV/c, K_S^0 , Λ and $\bar{\Lambda}$ show a pattern similar

to that observed in Au–Au collisions at top RHIC energy, although the R_{CP} values are found to be larger at SPS. At RHIC, this pattern has been interpreted in the framework of models that combine parton energy loss with hadronization via coalescence, at intermediate p_T , and via fragmentation, at higher p_T . The measured K_S^0 0–5%/40–55% R_{CP} is not reproduced by a theoretical calculation that includes only initial-state nuclear effects. The data can be better described by including final-state parton energy loss as predicted for SPS energy on the basis of RHIC data.

Acknowledgements

We thank V. Greco, X.N. Wang and U.A. Wiedemann for fruitful discussions and T. Peitzmann for making available to us the WA98 data.

References

- [1] S.S. Adler, et al., PHENIX Collaboration, Phys. Rev. C 69 (2004) 034910;
J. Adams, et al., STAR Collaboration, Phys. Rev. Lett. 91 (2003) 172302.
- [2] J. Adams, et al., STAR Collaboration, Phys. Rev. Lett. 92 (2004) 052302.
- [3] I. Arsene, et al., BRAHMS Collaboration, nucl-ex/0410020;
K. Adcox, et al., PHENIX Collaboration, nucl-ex/0410003;
B.B. Back, et al., PHOBOS Collaboration, nucl-ex/0410022;
J. Adams, et al., STAR Collaboration, nucl-ex/0501009.

- [4] M. Gyulassy, X.N. Wang, *Nucl. Phys. B* 420 (1994) 583;
M. Gyulassy, X.N. Wang, *Phys. Rev. Lett.* 68 (1992) 1480.
- [5] R. Baier, D. Schiff, B.G. Zakharov, *Annu. Rev. Nucl. Part. Sci.* 50 (2000) 37;
C.A. Salgado, U.A. Wiedemann, *Phys. Rev. D* 68 (2003) 014008.
- [6] D. d’Enterria, PHENIX Collaboration, *J. Phys. G* 31 (2005) S491;
J. Klay, STAR Collaboration, *J. Phys. G* 31 (2005) S451.
- [7] M.M. Aggarwal, et al., WA98 Collaboration, *Eur. Phys. J. C* 23 (2002) 225.
- [8] D. d’Enterria, *Phys. Lett. B* 596 (2004) 32.
- [9] S.A. Voloshin, *Nucl. Phys. A* 715 (2003) 379;
R.C. Hwa, C.B. Yang, *Phys. Rev. C* 67 (2003) 064902;
R.J. Fries, B. Muller, C. Nonaka, S.A. Bass, *Phys. Rev. C* 68 (2003) 044902;
V. Greco, C.M. Ko, P. Levai, *Phys. Rev. C* 68 (2003) 034904.
- [10] P.R. Sorensen, STAR Collaboration, *J. Phys. G* 31 (2005) S889.
- [11] V. Manzari, et al., NA57 Collaboration, *J. Phys. G* 25 (1999) 473.
- [12] F. Antinori, et al., NA57 Collaboration, *Eur. Phys. J. C* 18 (2000) 57.
- [13] F. Antinori, et al., NA57 Collaboration, *J. Phys. G* 31 (2005) 321.
- [14] R.J. Glauber, G. Matthiae, *Nucl. Phys. B* 21 (1970) 135;
A. Bialas, M. Bleszyński, W. Czyz, *Nucl. Phys. B* 111 (1976) 461.
- [15] A.S. Carroll, et al., *Phys. Lett. B* 61 (1976) 303.
- [16] R. Gottgens, et al., *Z. Phys. C* 19 (1983) 283.
- [17] C.W. de Jager, H. de Vries, C. de Vries, *At. Data Nucl. Data Tables* 14 (1974) 479.
- [18] E. Andersen, et al., WA97 Collaboration, *Phys. Lett. B* 433 (1998) 209;
R.A. Fini, et al., WA97 Collaboration, *J. Phys. G* 27 (2001) 375.
- [19] F. Antinori, et al., NA57 Collaboration, *J. Phys. G* 30 (2004) 823.
- [20] J. Podolansky, R. Armenteros, *Philos. Mag.* 45 (1954) 13.
- [21] F. Antinori, et al., NA57 Collaboration, *Phys. Lett. B* 595 (2004) 68.
- [22] S.S. Adler, et al., PHENIX Collaboration, *Phys. Rev. Lett.* 91 (2003) 172301.
- [23] G.E. Bruno, et al., NA57 Collaboration, *J. Phys. G* 30 (2004) S717.
- [24] D. Evans, et al., WA94 Collaboration, WA85 Collaboration, *J. Phys. G* 25 (1999) 209.
- [25] S.S. Adler, et al., PHENIX Collaboration, *Phys. Rev. C* 69 (2004) 034909.
- [26] J. Cronin, et al., *Phys. Rev. D* 11 (1975) 3105;
D. Antreasyan, et al., *Phys. Rev. D* 19 (1979) 764.
- [27] F. Matathias, et al., PHENIX Collaboration, *J. Phys. G* 30 (2004) S1113;
D. Kotchetkov, et al., PHENIX Collaboration, *J. Phys. G* 30 (2004) S1317.
- [28] K.J. Eskola, V.J. Kolhinen, C.A. Salgado, *Eur. Phys. J. C* 9 (1999) 61.
- [29] X.N. Wang, private communication.
- [30] X.N. Wang, *Phys. Rev. C* 61 (2000) 064910;
X.N. Wang, *Phys. Rev. Lett.* 81 (1998) 2655;
X.N. Wang, *Phys. Lett. B* 595 (2004) 165.
- [31] A. Dainese, C. Loizides, G. Paic, *Eur. Phys. J. C* 38 (2005) 461;
A. Dainese, C. Loizides, G. Paic, private communication.