



REVIEW

10.1002/2015JG003131

Special Section:

Arctic Freshwater Synthesis

Key Points:

- We review processes across hydrophysiographic regions of the terrestrial Arctic freshwater system
- Arctic hydrologic change affects atmosphere, ecology, resources, and oceans
- Interfaces between hydrology and other Earth system components are critical

Correspondence to:

A. Bring,
arvid.bring@unh.edu

Citation:

Bring, A., I. Fedorova, Y. Dibike, L. Hinzman, J. Mård, S. H. Mernild, T. Prowse, O. Semenova, S. L. Stuefer, and M.-K. Woo (2016), Arctic terrestrial hydrology: A synthesis of processes, regional effects, and research challenges, *J. Geophys. Res. Biogeosci.*, 121, 621–649, doi:10.1002/2015JG003131.

Received 1 JUL 2015

Accepted 2 FEB 2016

Accepted article online 5 FEB 2016

Published online 30 MAR 2016

©2016. The Authors.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Arctic terrestrial hydrology: A synthesis of processes, regional effects, and research challenges

A. Bring^{1,2,3}, I. Fedorova^{4,5}, Y. Dibike^{6,7}, L. Hinzman⁸, J. Mård⁹, S. H. Mernild^{10,11}, T. Prowse^{6,7}, O. Semenova^{5,12}, S. L. Stuefer¹³, and M.-K. Woo¹⁴

¹Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, Durham, New Hampshire, USA, ²Department of Physical Geography, Stockholm University, Stockholm, Sweden, ³Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden, ⁴Arctic and Antarctic Research Institute, St. Petersburg, Russia, ⁵Hydrology Department, Institute of Earth Sciences, St. Petersburg State University, St. Petersburg, Russia, ⁶Environment Canada, Victoria, British Columbia, Canada, ⁷Water and Climate Impacts Research Centre, University of Victoria, Victoria, British Columbia, Canada, ⁸University of Alaska Fairbanks, Fairbanks, Alaska, USA, ⁹Program for Air, Water and Landscape Sciences, Department of Earth Sciences, Uppsala University, Uppsala, Sweden, ¹⁰Antarctic and Subantarctic Program, Universidad de Magallanes, Punta Arenas, Chile, ¹¹Faculty of Engineering and Science, Sogn og Fjordane University College, Sogndal, Norway, ¹²State Hydrological Institute, St. Petersburg, Russia, ¹³Department of Civil and Environmental Engineering, College of Engineering and Mines, University of Alaska Fairbanks, Fairbanks, Alaska, USA, ¹⁴School of Geography and Earth Sciences, McMaster University, Hamilton, Ontario, Canada

Abstract Terrestrial hydrology is central to the Arctic system and its freshwater circulation. Water transport and water constituents vary, however, across a very diverse geography. In this paper, which is a component of the Arctic Freshwater Synthesis, we review the central freshwater processes in the terrestrial Arctic drainage and how they function and change across seven hydrophysiographical regions (Arctic tundra, boreal plains, shield, mountains, grasslands, glaciers/ice caps, and wetlands). We also highlight links between terrestrial hydrology and other components of the Arctic freshwater system. In terms of key processes, snow cover extent and duration is generally decreasing on a pan-Arctic scale, but snow depth is likely to increase in the Arctic tundra. Evapotranspiration will likely increase overall, but as it is coupled to shifts in landscape characteristics, regional changes are uncertain and may vary over time. Streamflow will generally increase with increasing precipitation, but high and low flows may decrease in some regions. Continued permafrost thaw will trigger hydrological change in multiple ways, particularly through increasing connectivity between groundwater and surface water and changing water storage in lakes and soils, which will influence exchange of moisture with the atmosphere. Other effects of hydrological change include increased risks to infrastructure and water resource planning, ecosystem shifts, and growing flows of water, nutrients, sediment, and carbon to the ocean. Coordinated efforts in monitoring, modeling, and processing studies at various scales are required to improve the understanding of change, in particular at the interfaces between hydrology, atmosphere, ecology, resources, and oceans.

1. Introduction

Many of the environmental changes currently under way in the Arctic involve terrestrial freshwater. Central examples include a decreasing extent and duration of snow cover [Callaghan *et al.*, 2011; Vaughan *et al.*, 2013], increasing flow from large Siberian rivers [Peterson *et al.*, 2002, 2006; McClelland *et al.*, 2006; Shiklomanov and Lammers, 2009; Overeem and Syvitski, 2010] and glaciers and ice sheets [Mernild and Liston, 2012; Church *et al.*, 2013], and changing partitioning between surface water and groundwater [L. C. Smith *et al.*, 2007; Walvoord and Striegl, 2007]. These changes illustrate that the Arctic Freshwater System also comprises feedbacks between the terrestrial hydrology and the ocean, atmosphere, ecosystems, and natural resources [Hinzman *et al.*, 2013; Raymond *et al.*, 2013].

Over the past few decades, investigation efforts into the Arctic Freshwater System have increased rapidly, and a large body of research, as well as a number of synthesis reports, has made major contributions to our knowledge. Nevertheless, our understanding of many of the changes remain incomplete, particularly with respect to the effects on and feedbacks to the ocean, atmosphere, ecosystems, and natural resources and also with respect to the dynamics with which the ongoing rapid changes will unfold across various regions of the pan-Arctic hydrological drainage basin.




Figure 1. Overview of regions. Schematic map (polar stereographic projection) illustrating a number of major hydrophysiographic regions in the terrestrial pan-Arctic drainage basin. The outer boundary of the basin is drawn to consider all areas potentially contributing to Arctic Ocean freshwater inflow, when also including ocean freshwater transport through the Bering Strait and the North Atlantic (indicated with arrows; see discussion on contributing area in *Prowse et al.* [2015a]). Although we emphasize that there is substantial variation within the regional boundaries outlined here, and many details that are excluded in this map, these large regions highlight a set of overarching landscape types across the Arctic drainage basin.

This paper is a part of the Arctic Freshwater Synthesis (AFS), a review effort initiated by the Climate and Cryosphere (CliC) project of the World Climate Research Program (WCRP), the International Arctic Science Committee (IASC), and the Arctic Monitoring and Assessment Program (AMAP), to provide updated information about the AFS. The AFS introduction paper [*Prowse et al.*, 2015a] contains background and can be consulted for a detailed discussion of the study domain, the pan-Arctic drainage basin. The AFS also comprises four other component papers that focus on freshwater processes related to the ocean [*Carmack et al.*, 2016], atmosphere [*Vihma et al.*, 2016], ecosystems [*Wrona et al.*, 2016], and natural resources [*Instanes et al.*, 2016], respectively. A fifth AFS component paper addresses modeling issues related to the AFS [*Lique et al.*, 2016], and a summary paper [*Prowse et al.*, 2015b] concludes the synthesis with key points and recommendations from the entire AFS.

In this AFS component paper, we provide a synthesis of terrestrial Arctic hydrology processes, their change drivers, and the main research challenges associated with them. Furthermore, a main aim is to identify linkages between terrestrial hydrology processes and other components of the AFS, and we therefore refer to other AFS papers throughout. We also attempt to separate distinctive aspects of freshwater processes, fluxes, and storages, and their changes, across representative hydrophysiographic regions of the AFS study domain, the pan-Arctic drainage basin (Figure 1), something that has received less focus in previous investigations (for brevity, the term Arctic is used in this paper to refer to the AFS domain).




Figure 2. Climatology of Arctic hydrophysiographical regions. Annual averages of grid-based (a) temperature, (b) precipitation, (c) evapotranspiration, and, as a proxy for runoff, (d) precipitation minus evapotranspiration values, over the regions defined in Figure 1 for the period 1979–2013. Regions are abbreviated as AT: Arctic tundra, BP: Boreal plains, S: Shields, M: Mountains, and G: Grasslands. Error bars denote one standard deviation of annual means, and open circles denote maximum and minimum annual means during the period. All data are from the ERA-Interim reanalysis product and are available at <http://apps.ecmwf.int/datasets/data/interim-full-daily>. Glaciers and ice sheets are not included here due to low density of observations in these ecoregions.

The hydrophysiographic regions we use in this paper mostly correspond to the terrestrial ecoregions defined by the World Wide Fund for Nature [Olson *et al.*, 2001]. The ecoregions (Arctic tundra, boreal plains, grasslands, and glaciers/ice caps) are used throughout the AFS, and are discussed more in detail in Prowse *et al.* [2015a]. However, due to the distinct hydrological and hydrogeological properties of mountains, shield regions, and wetlands, we also include those three categories as additional hydrophysiographic regions in this AFS component. The data sets used to define these regions are, for mountains, a study by Adam *et al.* [2006], for shield regions, the Canadian Geological Survey [Kirkham *et al.*, 1995], and for wetlands, the Global Lakes and Wetlands Database [Lehner and Döll, 2004]. We stress that each of these regions extends over a large area and therefore comprises quite diverse environments where local variations in hydrology are likely to be substantial. However, despite inevitable ambiguity and internal diversity, each region has still been identified as distinct in the aforementioned sources, and we use them here as overarching categories of hydrophysiographical landscape types.

2. System Functioning and Key Processes

This section presents a short description of the main processes pertaining to the Arctic terrestrial freshwater system, with focus on freshwater storages and fluxes. We discuss precipitation, evapotranspiration, surface runoff and channel flows, permafrost and groundwater hydrology, and river and lake ice. Following the processes, we highlight the central characteristics of a number of Arctic hydrophysiographical regions, also with regard to freshwater, its main storages, and fluxes. These regions are outlined in Figure 1, and an overview of regionally averaged temperatures and water fluxes based on gridded data is presented in Figure 2 (see also further discussions in Lique *et al.* [2016], Prowse *et al.* [2015a], and Vihma *et al.* [2016]).

Throughout the remainder of the paper, we progress from describing the system (section 2) to treating its past and future changes (sections 3–4). We then review key linkages across components of the Arctic freshwater system (section 5) and finally highlight research needs (section 6). As in this section, all of the following sections 3–6 are separated into a first section where we discuss processes (with some regional examples occasionally highlighted) and a second section where we review particular effects in each

hydrophysiographical region. With this organization, we hope that a reader interested in specific processes or regions is able to quickly find the right material in the paper.

2.1. Processes

Precipitation is the major flux into the terrestrial freshwater system (see *Lique et al.* [2016], *Prowse et al.* [2015a], and *Vihma et al.* [2016] for overviews and quantitative estimates). A substantial proportion of Arctic annual precipitation is falling and stored as snow, of heterogeneous spatial distribution [e.g., *Liston and Hiemstra*, 2011; *Mernild et al.*, 2014], and released to the river network in a relatively short window of time during spring snowmelt. Regionally, precipitation stored as snow contributes to runoff also in summer months. The phase of precipitation influences both annual and seasonal water balances at multiple spatial scales. The intensity with which precipitation is delivered also impacts runoff generation, with the spring freshet occasionally rivaled by summer thunderstorms for smaller basins [*Kane et al.*, 2008].

Over most Arctic basins, the majority of precipitation returns to the atmosphere as evapotranspiration (ET). ET links the water and energy cycles and couples the land to the atmosphere (evaporation over the Arctic Ocean is treated in more detail in *Vihma et al.* [2016]). The Arctic annual ET water flux is generally smaller than annual precipitation, except over a few southern inland areas [*Serreze et al.*, 2006, Figure 1] and over lakes and wetlands, where summer ET may exceed summer, or even annual, precipitation [*Marsh and Bigras*, 1988; *Rovansek et al.*, 1996; *Bowling and Lettenmaier*, 2010]. As the transpiration depends on the vegetation canopy, ET varies considerably even on local scales, as well as in time. The length of the snow cover and growing seasons are critical controls on the ET water flux. In addition, landscape variations such as lake area change may add up to considerable effects on ET and runoff [*Hinzman et al.*, 2005; *Karlsson et al.*, 2015]. Large-scale evapotranspiration values are difficult to estimate, but recent satellite-based assessments indicate pan-Arctic averages of $\sim 230 \text{ mm yr}^{-1}$ [*Zhang et al.*, 2009], with values ranging from 136 mm yr^{-1} in grasslands to 596 mm yr^{-1} in evergreen broadleaf forests [*Mu et al.*, 2009]. The ET data collected locally at eddy flux towers have been widely used for parameterization of hydrologic models [*Sun et al.*, 2008] and ET algorithms for remote sensing products [*Ruhoff et al.*, 2013]. Airborne eddy correlation measurements provide a useful means of bridging the scale interval between flux towers and regional models [*Sellers et al.*, 2012].

Besides ET, river discharge is the other major water flux out of Arctic basins. Freshwater flow through the Arctic's principal rivers conveys water, heat, sediments, carbon, and nutrients to the coastal domain and to the Arctic Ocean. Furthermore, along the river pathways, river flows, ice conditions, and runoff regimes control winter transportation, infrastructure and resource extraction, and ecosystem dynamics. Most of the water is transported to the Arctic Ocean during the spring snowmelt and in summer. The peak flow rates (May–June) can exceed the mean annual flow rate as much as 40 times for the Yenisey and Lena Rivers; the corresponding ratio for the Mackenzie River is much less, about 5 times [*Aagaard and Carmack*, 1989; *Bowling et al.*, 2000]. Due to dam construction, shifts of discharge strongly influence the seasonal flow for many Arctic rivers, including the Yenisey, Lena, and Mackenzie [*McClelland et al.*, 2004; *Yang et al.*, 2014]. In general, the variability is dampened by regulation, as spring and summer flow is held back and released in winter when flows are low. Effects on annual flows, however, are limited for most of the rivers [*McClelland et al.*, 2004; *Yang et al.*, 2004b; *Stuefer et al.*, 2011]. The largest Arctic rivers maintain channel flow year around including the winter flows under seasonal ice cover. Smaller northern rivers, however, often freeze to the bottom with no channel flow in winter. A portion of winter discharge is seasonally stored as river and lake ice and released in spring [*Prowse et al.*, 2011]. Formation and growth of afeis during winter are typical for many northern rivers [*Kane*, 1981]. In total, rivers deliver around $4 \times 10^3 \text{ km}^3$ of freshwater annually to the Arctic Ocean [*Serreze et al.*, 2006; *Haine et al.*, 2015], although this figure is strongly dependent on the total contributing area that is considered [*Prowse et al.*, 2015a].

Apart from the water itself, river conveyance of heat influences inland ice and ecosystem dynamics, particularly during transition seasons. The total delivery of water constituents such as nutrients, sediment, and carbon is less well known than that of the freshwater itself [*Bring and Destouni*, 2009], but recent estimates indicate fluxes of total nitrogen amounting to 1.3 Tg N yr^{-1} [*Holmes et al.*, 2011], total phosphorus 73 Gg P yr^{-1} [*Holmes et al.*, 2011], sediment $324\text{--}884 \text{ Tg yr}^{-1}$ [*Hasholt et al.*, 2006], inorganic carbon $57 \pm 9.9 \text{ Tg C yr}^{-1}$ [*Tank et al.*, 2012], and dissolved organic carbon $34\text{--}38 \text{ Tg C yr}^{-1}$ [*Holmes et al.*, 2011]. The water chemical composition is often strongly controlled by landscape and ecosystem processes [*Koch et al.*, 2013; *Aiken et al.*, 2014] (see *Wrona et al.* [2016], for further discussion). Carbon from rivers is a key input to near-coastal and ocean acidification, processes which are further discussed in *Carmack et al.* [2016].

Water chemistry and water flows in Arctic basins are both often influenced by permafrost. Active layer dynamics govern a wide range of surface and subsurface processes across permafrost landscapes and control mechanisms of runoff generation. The topographic relief, which varies between mountains, slopes, and flat terrain, is also strongly influencing soil moisture. *Carey and Woo* [2001], *Quinton et al.* [2005], and *Semenova et al.* [2013] have shown the impact of topography, soil and vegetation on ground freeze-thaw, soil moisture storage, surface and subsurface flow distribution, evaporation, and other processes. Due to the large extent of the area underlain by permafrost, the active layer thickness (ALT) and behavior varies across the Arctic, which influences soil moisture and storage. The mechanical stability of soil is also influenced by the water and ice content of the active layer [see *Instanes et al.*, 2016].

On shorter time scales, seasonal changes in surface ice are prominent characteristics of Arctic river systems. During winter, lake and river ice grow to cover $1.7 \times 10^6 \text{ km}^2$, an area approximately equal to the Greenland ice sheet. The peak volume of $1.6 \times 10^3 \text{ km}^3$ roughly matches the Northern Hemisphere snowpack on land [Brooks et al., 2013]. This freshwater ice produces numerous effects on physical systems, ecosystem services, and socioeconomic systems within Arctic freshwater storage and flow networks [Instanes et al., 2016; Wrona et al., 2016], although the hydrologic controls of many effects originate well outside sub-Arctic latitudes, via the headwaters of the large northward flowing rivers [Bennett and Prowse, 2010].

Most meteorological and climatological effects of variations in freshwater ice (e.g., coverage and duration) are confined primarily to the local or regional scale (e.g., radiation and convective fluxes), with the greatest effects produced by ice cover on large lakes [Rouse et al., 2005], although the process of river ice breakup has also been shown to be important, especially on the spring climate of large river deltas [Prowse et al., 2011]. However, effects on evapotranspiration, precipitation, and their feedbacks on hydrology may act on much larger scales, even influencing water balance of large Arctic basins [Rouse et al., 2008]. The magnitude and timing of hydrologic extremes such as low flows and floods are mostly controlled by the dynamics of river ice freezeup and breakup [Beltaos and Prowse, 2009], although reservoir discharge and groundwater base flow also control winter low flows [Woo and Thorne, 2014]. Spring breakup tends to be the dominant hydrologic event across the full domain of large rivers and deltas, such as the Mackenzie and the Lena [de Rham et al., 2008; Goulding et al., 2009; Fedorova et al., 2015], and also the main agent of sediment transport and morphological change [Turcotte et al., 2011].

2.2. Hydrophysiographic Regions

Here we summarize hydrological characteristics across the seven hydrophysiographic regions shown in Figure 1: Arctic tundra, boreal plains, shields, mountains, grasslands, glaciers and ice sheets, and wetlands. We also highlight a number of ways in which the hydrological responses to environmental changes can be expected to differ across the regions.

In the tundra, continuous permafrost, with an active layer that decreases from about 1.5 m in the south to 0.5 m or less in the north (Circumpolar Active Layer Monitoring data, available at <http://www.gwu.edu/~calm/data/north.html>), strongly influences water fluxes and storage. With limited permeability of the frozen soil (both seasonal and perennial) that restricts infiltration and water storage, groundwater storage and circulation are largely confined to the seasonally thawed zone. Generally, watersheds with a high percentage of permafrost coverage have low subsurface storage capacity for liquid water and thus a low winter base flow (minimum flow) and a high spring or summer peak flow (maximum flow) [Woo, 1986; Kane, 1997; Yang et al., 2002; L. C. Smith et al., 2007; Ye et al., 2009]. Evapotranspiration and precipitation are both very low, although evapotranspiration can be highly variable in shrublands [Mu et al., 2009]. At its northernmost extreme, the tundra is adjacent to the ocean, where groundwater and smaller streams transport relatively poorly quantified amounts of water, nutrients, sediment, and carbon to the ocean.

In boreal plains, the low gradients lead to slow surface and subsurface water movement and ample depression storage, which is associated with the extensive occurrence of wetlands [Ferone and Devito, 2004]. Permafrost becomes discontinuous in the south, which allows groundwater to circulate more freely in relation to the impervious frozen substrate in the southern boreal plains [K. B. Smith et al., 2007]. With increasing solar radiation and vegetation cover, evapotranspiration flux increases southward [Mu et al., 2009]. Seasonal storage of snow is less pronounced than in the tundra, and high flow events arising from rain become more pronounced southward, next to the spring freshet [Su et al., 2005].

In shield regions, the bedrock matrix is essentially impermeable but may be riddled with joints and fractures that, depending on their content, permit infiltration [Spence and Woo, 2002]. Thus, the runoff ratio can vary greatly on small scales. The major structural fissures are eroded into soil-filled valleys with lakes and wetlands that form tortuous drainage networks [Spence and Woo, 2003]. Groundwater yield is low relative to surface water, as permafrost and bedrock allow limited groundwater storage. Instead, surface storage in lakes and wetlands is central to runoff generation, which follows the principle of fill and spill, whereby lakes and wetlands in valleys have to be filled above the elevation of their outlet levels before flow commences [Woo and Mielko, 2007]. Otherwise, water is held back in storage and discharge is interrupted, leading to cessation of flow downstream.

Mountainous regions are responsible for >60% of the annual flow of Mackenzie Basin [Woo and Thorne, 2003] and are important freshwater sources also for larger basins in eastern Siberia and the Russian Far East. The Lena River receives on average 40% of the annual flow from the mountainous regions of the Aldan and Upper Lena Rivers [Berezovskaya et al., 2005]. The large altitudinal range of mountainous regions leads to a prominent vertical zonation and an aspect-controlled microclimate. In high elevations, and at high latitudes also at lower elevation, water is stored interannually in snowpack and glaciers, with gradual melt release prolonging the duration of flow generation over the snow-free season. Orographic effects generally yield high precipitation, particularly in coastal regions [Mernild et al., 2015]. Aspect plays an important role in the energy and water balances of slopes and tributary basins. Snowmelt on south facing slopes can be a month in advance of north facing slopes, and evapotranspiration is higher on south than north slopes. For example, for Wolf Creek, Yukon, evapotranspiration on a north slope averaged 315 mm yr^{-1} and on the opposite slope it reached 372 mm yr^{-1} [Carey and Woo, 2001].

Some large Arctic rivers pass through prairies (e.g., the Saskatchewan River that joins the Nelson River) and steppes (e.g., the Ob) in their upper courses. These low-relief areas allow air masses and their accompanying disturbances to sweep uninterrupted over great distances. This leads to large climatic variability that impacts terrestrial hydrology, principally with regard to floods and droughts. In addition to flooding from convective rainfall, mountain rivers also effectively deliver snowmelt floods to the plains. In summer, low precipitation and high temperatures combine to effect large net water loss to evaporation. However, evaporation can be very variable due to large fluctuations in climate and surface conditions [Armstrong et al., 2015]. Within this region, some basins have no outlet and form hydrological enclaves, disconnected from the surrounding watershed. During dry years, the low precipitation and decrease in snowpack lead to drying out of local wetlands [Fang and Pomeroy, 2008], which then become disconnected from each other [Shaw et al., 2012].

Glaciers and ice sheets play a major role in the Arctic freshwater system through their impact on the surface energy budget, the water cycle, and sea level [Vaughan et al., 2013]. Freshwater discharge from the Greenland ice sheet and the Arctic terrestrial rivers, the latter substantially larger, has a critical influence on Arctic Ocean circulation [see Carmack et al., 2016; Vihma et al., 2016; Lique et al., 2016]. For smaller Arctic glaciers and ice caps, their role in freshwater storage and flux varies greatly throughout the region. With the exception of the Yukon River basin, glaciers contribute only a small share of flow for all the major Arctic rivers [Dyurgerov et al., 2010]. Regionally, however, they strongly affect both runoff seasonality and water storage change, as freshwater flow is redistributed to the summer.

Wetlands are prominent through the Arctic, especially in the tundra, where they can function as large systems of lakes and wetlands, with water accumulating in depressions. There are different kinds of wetlands in the Arctic, with a central distinction differentiating between peatlands and wetlands in the general sense. In boreal western Siberia and Canada, peatlands are more extensive, often with a water-storing peat layer 1 to 5 m thick [Sheng et al., 2004], whereas wetlands in the tundra generally do not have thick peat layers. Along watercourses, wetlands tend to reduce the variability in river flows, both storing water during high flows and acting as reservoirs of runoff generation during dry summer periods. In the tundra, small lakes sometimes form systems where large amounts of water may accumulate, with high propensity for thermokarst development. When thawing, ice complexes in Yedoma soils form secondary thermokarst objects called *alases* [Morgenstern et al., 2011]. Wetlands and peatlands have high adsorption and ion exchange capacity, which may lead to accumulation of water-transported metals (including heavy metals) and carbon. Wetlands also contribute to retention of pollutants and nutrients, although this depends on the share of river flow that circulates through the wetland [Quin et al., 2015]. The ion composition and suspended material flows differ during summer flows and high flows [Raymond et al., 2007; Frey and McClelland, 2009] (see also Wrona et al. [2016], for further discussion).

In addition to the regions described above, the riverine coastal domain [see *Carmack et al.*, 2016] could be separately considered as a region with unique hydrodynamic and biogeochemical properties, which may also grow in importance as erosion progresses, permafrost thaws, and sea ice declines. Along the coast, erosion, tidal, and surge processes, as well as water, heat, and geochemical fluxes, are under joint influence of riverine, marine, and atmospheric processes and affect riparian ecosystems and mesoscale coastal landscapes. This emerging research domain is discussed further in *Carmack et al.* [2016] and *Prowse et al.* [2015b].

3. Past Changes and Key Drivers

In this section, we review past changes to the main processes of the Arctic terrestrial freshwater system, with focus on freshwater storages and fluxes. Following the processes, we also highlight past changes in the Arctic hydrophysiographical regions outlined in Figure 1.

3.1. Processes

With regard to precipitation processes, we focus on snow cover changes (see *Vihma et al.* [2016] and *Lique et al.* [2016] for other changes to precipitation). Syntheses of several ground and satellite observational data sets indicate that Northern Hemisphere snow cover extent decreased by 2.2% per decade, averaged for March and April, and by 14.8% per decade for June, over the period 1979–2012 [*Brown and Robinson*, 2011] (updated in *Vaughan et al.* [2013]). Both positive and negative regional trends are distributed throughout the pan-Arctic, however, including spatially distinct areas of increasing and decreasing snow water equivalent (SWE) or snow season length [*Callaghan et al.*, 2011]. In spite of strong regional variability—for example, increasing SWE in northern Eurasia—snow is, by most measures, generally decreasing throughout the Arctic as shown by *Liston and Hiemstra* [2011]. They used a physically based spatially distributed snow modeling system (SnowModel) in conjunction with NASA Modern-Era Retrospective Analysis for Research and Applications atmospheric reanalysis data to show that snow cover onset is later, the snow-free date in spring arrives earlier, and snow cover duration has decreased over the period 1979–2009 [*Liston and Hiemstra*, 2011]. These changes in snowpack are related to the earlier onset of the snowmelt runoff [*Tan et al.*, 2011]. Despite the shorter duration of snow cover, however, hydrological modeling experiments show that an increase in SWE, as observed in the northern parts of the large Eurasian basins, may still lead to increased annual runoff [*Troy et al.*, 2012]. Main drivers of snow changes are air temperature increase and changing amounts and timing of precipitation. The modeled sensitivity of snow to these drivers varies strongly across climates and regions [*Brown and Mote*, 2009]. As winter air temperature increases, observations of rain-on-snow events become more common in Arctic regions where they were rarely seen before [*Ye et al.*, 2008; *Nowak and Hodson*, 2013; see also *Vihma et al.*, 2016; *Instanes et al.*, 2016; *Wrona et al.*, 2016].

In terms of evapotranspiration, it is challenging to determine large-scale changes as observations are scarce. *Zhang et al.* [2009] developed an ET algorithm driven by satellite remote sensing inputs to assess spatial patterns and temporal trends in ET over the pan-Arctic basin 1983 to 2005 and found a mean trend of $+0.38 \text{ mm yr}^{-2}$ ($p < 0.05$). Negative ET trends occurred over 32% of the region, primarily in the boreal forests of southern and central Canada. *Rawlins et al.* [2010] showed that model trends of annual ET are significantly positive, with a multimodel mean trend (1950–1999) of $+0.17 \text{ mm yr}^{-2}$ ($p < 0.1$). The differences between the estimates suggest an amplification of ET increases over the recent decades. Increased ET is to be expected with observed warming due to the ability of the atmosphere to hold more moisture at higher temperatures, but ET changes are also driven by landscape alterations, such as vegetation shifts, fires, and lake and reservoir changes [*Wrona et al.*, 2016]. For example, increases in shrubs and trees enhance evapotranspiration and thereby the loss of water from the basin, which leads to drier surface conditions [*Wrona et al.*, 2016].

A large-scale pattern of observed river discharge increases has been frequently reported for Arctic rivers [*Peterson et al.*, 2002, 2006; *McClelland et al.*, 2006; *Shiklomanov and Lammers*, 2009; *Dyurgerov et al.*, 2010; *Overeem and Syvitski*, 2010; *Holmes et al.*, 2013; *Bring and Destouni*, 2014], although decreases are also noted, particularly for some North American rivers [*Déry et al.*, 2005; *McClelland et al.*, 2006]. However, over the last couple of decades, discharge in North American rivers reversed this trend and have increased as well [*Déry et al.*, 2009], aligning them with the more general pattern of increasing discharge. The total river flows, calculated from averages of reanalysis and in situ data, have likely increased from $3900 \pm 390 \text{ km}^3$ during 1980–2000 to $4200 \pm 420 \text{ km}^3$ during 2000–2010 [*Haine et al.*, 2015], with errors assumed to be about 10%. In addition to average flows, changes in low and high flows have also been reported. *Spence et al.* [2014] have shown that

increased late autumn rains may cause the enhancement of winter flow and impact geochemical fluxes from headwater catchments of the subarctic Canadian Shield. *Shiklomanov et al.* [2007] reported that increases and decreases in station maximum daily flows were equally common, with very few significant trends in summer months, which led them to question the generally expected future increase in floods. Increases in station low flows, which are reported for several regions [*L. C. Smith et al.*, 2007; *St. Jacques and Sauchyn*, 2009; *Ehsanzadeh and Adamowski*, 2010; *Rennermalm et al.*, 2010; *Karlsson et al.*, 2012; *Walvoord et al.*, 2012; *Karlsson et al.*, 2015], are consistent with recession flow analyses [*Lyon et al.*, 2009; *Lyon and Destouni*, 2010; *Brutsaert and Hiyama*, 2012] and modeling of permafrost thaw [*Bense et al.*, 2009, 2012; *Frampton et al.*, 2011, 2013] that indicate an increasing contribution of groundwater to streamflow [*Walvoord and Striegl*, 2007]. Overall, a number of reanalysis, modeling, and observation-based studies show that increased atmospheric moisture transport (AMT), due to higher temperatures and changing atmospheric circulation, is likely the principal driver of long-term increases in flow [*Zhang et al.*, 2008, 2013; *Rawlins et al.*, 2009; *Troy et al.*, 2012] (see discussion of atmospheric changes in *Vihma et al.* [2016]). Model-observation studies have also shown that winter precipitation stored as snow is a central component of the AMT contribution [*Troy et al.*, 2012; *Zhang et al.*, 2013]. In contrast, ground ice melt and evaporation from increasingly ice-free ocean water have not been responsible for any greater contribution to historical flow increases [*McClelland et al.*, 2004; *Pavelsky and Smith*, 2006; *Zhang et al.*, 2013], although an exact quantification remains elusive.

Permafrost thaw has changed the hydrological regime in some basins, particularly through altered surface and subsurface interactions. For example, *Connon et al.* [2014] report observed increases of annual runoff in the lower Liard Valley (NWT, Canada) by between 112 and 160 mm over the period of 1996–2012, mainly due to increase of plateau runoff contributing areas and a change in the relative proportions of the major land cover types, such as peat plateaus, channel fens, and flat bogs. These values are large relative to average annual runoff values of between 147 and 216 mm for the period. Changes to temperature and precipitation have also interacted to produce changes in evapotranspiration, runoff, peak seasonal snow accumulation, and snow season length in permafrost basins. Short-term changes in air temperature, ice cover, and soil moisture do not trigger systematic hydrological shifts in permafrost, although they provide a memory that manifests itself during the next warm season, as shown in a model-observation study [*Park et al.*, 2013]. Long-term changes in climate, however, alter the landscape structure and its runoff formation, principally through an increased ALT, as indicated by both observations and modeling [*Quinton et al.*, 2011; *Frampton et al.*, 2013]. In Eurasia, the ALT has generally increased due to thicker snowpacks and high summer soil moisture [*Park et al.*, 2013]. In the Mackenzie and Yukon basin, on the other hand, combined effects of less insulation caused by thinner snow depth and drier soil during summer have partly offset warming effects on the ALT [*Park et al.*, 2013], although these results are based on modeling and further research is needed to confirm them.

Ice cover observations on Northern Hemisphere lakes indicate a shortening duration, with the time of breakup generally changing more rapidly than the freezeup [*Benson et al.*, 2011]. Trends over 1855–2004 were steeper than over 1905–2004, but the most rapid changes occurred in the most recent 30 year period, with freezeup 1.6 d/decade later, breakup 1.9 d/decade earlier, and ice duration 4.3 d/decade shorter. Although ice cover tends to be more sensitive to air temperature variations at lower than at higher latitudes [*Livingstone et al.*, 2010], remote sensing observations indicate that ice cover loss seems to be more rapid in very high latitude lakes [*Latifovic and Pouliot*, 2007]. High-latitude lakes lost ice cover at 1.75 d yr^{-1} during the 1970–2004 period of most rapid depletion, which is more than 4.5 times the rate of lakes in southern Canada. It is unclear whether this reflects the more recent and greater high-latitude warming or potential differences in observational techniques [*Prowse and Brown*, 2010]. Potential effects of reduced lake ice cover on feedbacks to the atmosphere are further discussed in *Vihma et al.* [2016].

In terms of river ice observations, *Beltaos and Prowse* [2009] noted an almost universal trend toward earlier breakup dates but considerable spatial variability in those for freezeup. Changes are often more pronounced during the last few decades of the twentieth century. Overall, twentieth century warming has led to a 10 to 15 day advance in breakup and delay in freezeup, respectively [see also *Lique et al.*, 2016]. This is in agreement with earlier estimates [*Magnuson et al.*, 2000], but the relationship is complicated by changes in snow accumulation and spring runoff [*Beltaos and Prowse*, 2009].

The discharge from large Arctic rivers (examples from Eurasian rivers shown in Figure 3) integrates all features and change drivers from the hydrophysiographic regions that make up their catchments [*Holmes et al.*, 2013]. The seasonal character of discharge, which for many large Arctic rivers is affected by dams [*McClelland et al.*,

2004; Yang *et al.*, 2014], shows quite high flow during all of the summer period and very low flows in winter (Figure 3a). Interannual variability is high, and the total water flow in a wet water year can be twice the flow in a dry year (Figure 3b). As noted above, there are examples of both increases and decreases in river discharge to the Arctic Ocean (Figure 3c), but for the Arctic as a whole, flow has increased both from Eurasia and North America when considering the longest possible time periods (NOAA Arctic report card 2011, available at www.arctic.noaa.gov/reportcard [Holmes *et al.*, 2013]).

3.2. Hydrophysiographic Regions

In this section, past changes and key drivers are summarized for each hydrophysiographic region (Figure 1). Annual river flows observed in the tundra region have generally increased, but there are exceptions [Zhang *et al.*, 2009; Overeem and Syvitski, 2010]. Increasing temperatures, and to a large degree, precipitation, are the main drivers of change in Arctic tundra hydrology [see Vihma *et al.*, 2016; Lique *et al.*, 2016]. The warming effect in sheltered locations has created polar oases in the otherwise barren high Arctic [Edlund and Alt, 1989], and remote sensing shows that recent warming has led to increased development of thermokarst lakes in ice-rich permafrost environments [Smith *et al.*, 2005]. Consecutive positive observed anomalies of snow depth and rainfall have likely contributed to the widespread warming of near-surface permafrost in the central Lena River basin [Iijima *et al.*, 2010]. Satellite estimates of evapotranspiration in the tundra show increases by 2.7–2.8 mm/decade over 1983–2005, but this figure is small in relation to annual values and only significant at $p=0.1$ for North America, and not significant for Eurasia [Zhang *et al.*, 2009; see also Vihma *et al.*, 2016].

Shrub expansion has occurred throughout the Arctic tundra during the past 50 years [Sturm *et al.*, 2001, 2005; Tape *et al.*, 2006, 2012], modifying the microclimate of the landscape and altering erosion and biogeochemical fluxes. The underlying mechanisms of shrubification are largely attributed to increasing air and soil temperature, but factors related to local hydrological changes such as precipitation, soil moisture, and snowpack also influence shrub growth [Wrona *et al.*, 2016].

While there is evidence of northward advance of the tree line across the boreal plains, with increase in tree density and canopy size of individual trees [Danby and Hik, 2007], other parts of the boreal forest are experiencing degradation of their discontinuous permafrost substrate, which has reduced the occurrence of tall-growing vegetation [Quinton *et al.*, 2011]. Large-scale satellite-based estimates indicate that evapotranspiration has increased in Eurasian forests (6.94 mm/decade, $p < 0.05$) but that it may have decreased in North American ones (−3.06 mm/decade, no significance) [Zhang *et al.*, 2009]. River flows have generally increased in the boreal plains and shield regions, but decreases are reported for some Canadian rivers [Zhang *et al.*, 2009; Déry *et al.*, 2011].

In shield regions, extensive hydropower development [see Instanes *et al.*, 2016] has also modified the flow of many rivers, lowering summer flows and increasing winter flows [Déry *et al.*, 2011]. Information on undeveloped rivers, with flow patterns unaltered by dam construction, is relatively scarce in the Arctic shield regions. For the Tana River in northern Norway and Finland, long-term station records do not reveal any significant trends in streamflow but indicate increases in both annual precipitation and annual precipitation variability [Dankers, 2003]. In contrast, streamflow variability has decreased in unregulated rivers on the Canadian Shield that drain into Hudson Bay [Déry *et al.*, 2011].

In some mountainous regions, thawing of alpine permafrost has modified the pathway, timing, and amount of runoff. Using measured data and reinforced by modeling based on Frampton *et al.* [2013], Sjöberg *et al.* [2013] found that accompanying observed permafrost degradation in northern Sweden, there are higher recession flows and increased winter discharge sustained by groundwater in most alpine basins they studied, although these signals were not consistent across landscapes and, for minimum flows, not correlated strongly with permafrost extent. Mountainous regions also contain intermontane basins. Compared with their surrounding heights, these basins have lower elevations and gentler topography and show earlier response to climatic change. In the Yukon Flats in Alaska, there are indications from remote sensing that expanded shallow supra-permafrost or intrapermafrost taliks are linked to the expansion and shrinking of lakes [Jepsen *et al.*, 2013].

On the low-latitude grasslands in the upstream portion of the largest Arctic river basins, reservoirs of different sizes have been constructed for irrigation, increasing evaporation. Groundwater has also been exploited to water crops. In the upper Ob, water flows have decreased in summer and increased in winter, as a result of increasing water abstraction for agriculture and industry and flow regulation from dams [Yang *et al.*, 2004a]. Similar effects have been observed in the upper Yenisey basin [Yang *et al.*, 2004b].




Figure 3. Examples of discharge patterns from large Eurasian Arctic rivers. (a) Monthly average discharge for the combined Eurasian Arctic drainage. (b) Differences in discharge between an average, a dry, and a wet year for the Lena River. (c) The long-term annual river discharge for the six largest Eurasian Arctic rivers from 1936 to 2002. Due to the declining accuracy of gauge readings in Russia [McClelland *et al.*, 2015], we only include data until 2002 in Figure 3. Even considering gauge uncertainty in the period since then, flows distinctively reached a record high in 2007 [Rawlins *et al.*, 2009; Shiklomanov and Lammers, 2009], and the years from 2000 to at least 2009 have been wetter than the average for the period 1950–2009 [Walsh *et al.*, 2011]. All river data are available from the Global Runoff Data Centre (<http://www.bafg.de>) and the R-ArcticNET database (<http://www.r-arcticnet.sr.unh.edu>).

Of the several thousands of glaciers that are located in the pan-Arctic, only ~25–30 have ongoing operational mass balance programs to quantify glacier mass balance conditions and change [World Glacier Monitoring Service, 2013] (www.wgms.ch). Analyses of glacier area fluctuations, based on historical accounts, aerial photography, and satellite images, are more numerous. Since the early 1990s, the increasing glacier [e.g., Kaser *et al.*, 2006; Cogley, 2012; Vaughan *et al.*, 2013; Mernild *et al.*, 2014] and Greenland ice sheet (GrIS) net mass loss and surface runoff have followed atmospheric warming [e.g., Hanna *et al.*, 2008; Box and Colgan, 2013; Church *et al.*, 2013; see also Carmack *et al.*, 2016; Lique *et al.*, 2016]. Mass loss from the GrIS has increased rapidly, with recent estimates of $375 \pm 24 \text{ km}^3 \text{ yr}^{-1}$ for 2011–2014 [Helm *et al.*, 2014] (based on estimations from CryoSat-2) and $575 \pm 95 \text{ km}^3 \text{ yr}^{-1}$ for September 2011 to September 2012 (NOAA Arctic report card 2012, available at www.arctic.noaa.gov/reportcard and based on Gravity Recovery and Climate Experiment, a satellite mission used to detect changes in water mass). For Greenland in its entirety, runoff was recently estimated to $481 \pm 85 \text{ km}^3 \text{ yr}^{-1}$ for 1960–2010 [Mernild and Liston, 2012] (based on modeling). On the pan-Arctic scale, the cumulative glacier net mass balance has been negative for all glacier regions for the period 1979–2009 [Mernild *et al.*, 2014], and overall contributions to sea level change are positive (see glacier change summary in Table 1). For some smaller basins, mass balance change constitutes a substantial share of river runoff. For example, about 35% of runoff from the Mittivakkat Glacier basin in southeast Greenland originates from net mass balance loss [Liston and Mernild, 2012].

Studies of past changes to wetlands indicate both increases and decreases in areal extent. Generally, the changes in number and area of wetland lakes and ponds depend on the local permafrost and hydrological conditions. Lakes have mostly decreased in size and number in areas where discontinuous and sporadic permafrost thaw is in progress [Andresen and Loughheed, 2015]. In these areas, flow paths and water tables are becoming deeper, and lakes may also drain as the permafrost substrate is breached. However, other factors,

Table 1. Regional Examples of Glacier Area Changes and Sea Level Contributions in the Pan-Arctic Drainage Basin

Glacial Area Changes			
Region	Time Period	Change in Area	Reference
Yukon Territory	1958/1960–2006/2008	–22%	<i>Barrand and Sharp</i> [2010]
British Columbia	1985–2005	–11%	<i>Bolch et al.</i> [2010]
Interior northern Baffin Island	1958–2005	–55%	<i>Anderson et al.</i> [2008]
Southeast Baffin Island	1920–2000	–13%	<i>Paul and Svoboda</i> [2010]
Southeast Greenland	1986–2011	–27%	<i>Mernild et al.</i> [2012]
Northern Polar Urals	1953/1960–2000	–22%	<i>Shahgedanova et al.</i> [2012]
Suntar Khayata Region, North Asia	1945–2002/2003	–19%	<i>Ananicheva et al.</i> [2006]
Koryak Upland near Kamchatka Peninsula	1950s–2003	–67%	<i>Ananicheva and Kapustin</i> [2010]
Sea Level Contributions			
Region	Time Period	Contribution to Sea Level Equivalent (SLE) in mm yr ^{–1}	Reference
Alaska	1999–2009	0.13 ± 0.05	<i>Mernild et al.</i> [2014]
Arctic Canada (North)	1999–2009	0.10 ± 0.01	<i>Mernild et al.</i> [2014]
Greenland (nonice sheet)	1999–2009	0.09 ± 0.01	<i>Mernild et al.</i> [2014]

such as changes in temperature and evapotranspiration, are also driving forces [Karlsson et al., 2015]. In contrast, increases, decreases, or no change in lake size and number are reported from areas in continuous permafrost [Jones et al., 2011; Andresen and Lougheed, 2015].

4. Projected Changes and Key Drivers

In this section, we review main projected changes to the central processes of the Arctic terrestrial freshwater system, with focus on freshwater storages and fluxes. Following the processes, we also highlight projected hydrological changes across the Arctic hydrophysiographical regions outlined in Figure 1.

4.1. Processes

In Table 2, we synthesize reported projections for hydrological processes across the hydrophysiographical regions. Projected changes in seasonal snowfall and snow water equivalent are spatially variable and depend on local climate conditions. In very cold regions, increased winter precipitation will lead to a deeper snow cover (and higher SWE), while in warmer regions, higher temperatures will lead to the opposite [Räisänen, 2008; see also Lique et al., 2016]. However, other snow-related variables, such as snow cover extent (SCE), exhibit a more direct relationship with air temperature.

Brutel-Vuilmet et al. [2013] found a strong negative correlation between Northern Hemisphere spring SCE and the corresponding mean surface air temperature, and by 2080–2099, the average reduction of seasonal SCE varies from 7.2 ± 3.8% for RCP2.6 to 24.7 ± 7.4% for RCP8.5, relative to a 1986–2005 reference period. Other modeling studies have also showed increases in snowfall and SWE associated with the projected increase in cold season precipitation in northeastern Eurasia and northern Canada, while they decreased in more southerly places in which warming effects dominated [Räisänen, 2008; Deser et al., 2010; Krasting et al., 2013]. In these studies, the –10°C and –20°C winter air temperature isotherm lines represent transition boundaries for snowfall [Deser et al., 2010; Krasting et al., 2013] and SWE [Räisänen, 2008], respectively.

There is growing evidence for an increase in precipitation extremes, with relative increases generally exceeding those for annual mean precipitation under the projected 21st century global warming [Kharin et al., 2013; Sillmann et al., 2013; Vihma et al., 2016]. A statistically significant signal of increasing summer season precipitation across the Arctic as a whole is likely to emerge from the variability when global temperature rise surpasses 1.4°C, which is likely to occur around 2040 [Mahlstein et al., 2012]. Signals in extreme precipitation, however, as well as regional signals, may emerge earlier or later, and because models underestimate past changes, the time of emergence for the entire Arctic is likely too conservative [Mahlstein et al., 2012].

A diagnostic study of future evapotranspiration changes projected in CMIP5 climate models under the RCP45 scenario by Laine et al. [2014] shows a change on the order of 0.05 mm d^{–1} in winter and 0.25 mm d^{–1} in summer over a 100-year period for the terrestrial pan-Arctic (1980–2000 versus 2080–2100) [see also Lique et al., 2016].

Table 2. Projected Changes to Hydrological Processes Across Arctic Hydrophysiographical Regions

Process	Hydrophysiographical Region	Projected Changes	Specific Effects/Comments
Precipitation	Arctic tundra	Generally increasing annual precipitation (highest increase in winter and fall)	Increase in snow water equivalent, earlier snowmelt, decrease in seasonal snow cover period, warming of permafrost, and increase in coastal erosion
	Boreal plains	Generally increasing annual precipitation (highest increase in winter and fall), with some regional decreases in summer precipitation	Increase in soil moisture and runoff, midwinter snowmelt events and more frequent rain on snow events, possible soil moisture decrease (drying) in summer, decrease in maximum snow water equivalent, and decrease in seasonal snow cover extent in May/June
	Mountains	Generally increasing annual precipitation (highest increase in winter and fall), with some regional decreases in summer precipitation	Increase in soil moisture and runoff and decrease in seasonal snow cover extent in May/June
	Shields	Generally increasing annual precipitation (highest increase in winter and fall), with some regional decreases in summer precipitation	Increase in soil moisture and runoff, midwinter snowmelt events and more frequent rain on snow events, and possible soil moisture decrease (drying) in summer
	Ice sheets, ice caps, and glaciers	Increasing annual precipitation	Increase in snowpack/accumulation
	Wetlands	Increasing annual precipitation (highest increase in winter and fall)	Midwinter snowmelt events and more frequent rain on snow events and decrease in seasonal snow cover extent in May/June
	Grasslands	Small increase in annual precipitation (with decreases in summer precipitation)	Decreases in summer precipitation with possible increase in summer drought events
	Vegetation and ecosystem changes		
Evapotranspiration	Arctic tundra	Increasing annual evapotranspiration (more in summer than winter)	Possible increase in summer drought events
	Boreal plains	Increasing annual evapotranspiration (more in summer than winter)	Possible drying (decrease in soil moisture) in summer
	Mountains	Increasing annual evapotranspiration (more in summer than winter)	Possible drying (decrease in soil moisture) in summer
	Shields	Increasing annual evapotranspiration (more in summer than winter)	Possible drying (decrease in soil moisture) in summer
	Ice sheets, ice caps, and glaciers	Increasing annual evapotranspiration and sublimation	Increasing surface mass loss
	Wetlands	Increasing annual evapotranspiration (more in summer than winter)	Possible decrease in wetland area
	Grasslands	Increasing annual evapotranspiration (possible decrease in summer)	Possible decrease in summer evaporation due to decrease in summer precipitation
Runoff and river flux	Arctic tundra	General increase in mean, high, and low flows and limited areas of decrease in high and low flows	Possible high and low flow decreases in western Siberia
	Boreal plains	General increase in mean flows but limited areas of decrease and variable changes in high and low flows	Possible high and low flow decreases in western Siberia, southern Canada
	Mountains	Consistent increase in mean flows, high and low flows generally increasing	Possible low flow decreases in western Siberia, southern Canada
	Shields	Smaller increase in mean flows and variable changes in high and low flows	Possible low flow decreases in southern Canada and Scandinavia
	Ice sheets, ice caps, and glaciers	Increase in runoff as glacier mass loss increases, followed by eventual runoff decrease as glacier shrinkage counteracts the increased melting	Initially increasing surface ablation/mass loss, thereafter a drop in surface ablation following glacier shrinkage
	Wetlands	General increase in mean flows and variable changes in high and low flows	Possible high and low flow decreases in western Siberia and western Canada
	Grasslands	Variable change in mean, low, and high flows	Increases more likely in Asia and decreases more likely in North America
	Vegetation changes		
Permafrost and groundwater hydrology	Arctic tundra	Increases in active layer depth and additional degradation of permafrost will provide greater reservoirs for	Soil moisture increasing on plateaus. Vegetation changes will increase evapotranspiration. Succession of

Table 2. (continued)

Process	Hydrophysiographical Region	Projected Changes	Specific Effects/Comments
		subsurface storage of groundwater and near-surface soil moisture. General increase in low flows due to strengthened connections between surface water and groundwater and increase in runoff and erosion.	wetting and drying of landscapes. Suprapermafrost groundwater will increase, providing a limited reservoir for later season flow to streams and rivers.
	Boreal plains	Complete disappearance of permafrost in marginal areas of sporadic occurrence. General increase in low flows due to strengthened connections between surface water and groundwater.	Where groundwater table is below the surface water, infiltration may leave the surface drier. In areas of groundwater upwelling, increased connectivity may support wetland development.
	Mountains	Limited change in groundwater dynamics but possible increases in low flows for valley and wetland regions in areas of degrading permafrost	In regions of discontinuous permafrost, springs tend to be found on slopes where percolating water first encounters permafrost and water is forced to the surface. Location of springs may shift.
	Shields	Limited change in groundwater dynamics. In regions of increasing summer aridity, possible increasing seasonal loss of connectivity between lakes, wetlands, and rivers.	Zones of recharge and discharge over thick shields are mostly tied to fractures and will generally remain unchanged.
	Ice sheets, ice caps, and glaciers	Limited change in groundwater dynamics; however, increase in groundwater storage is possible.	Possible increase in groundwater storage if the increased rate of ice melt is greater than the rate of groundwater discharge. If groundwater reservoir already saturated (as is typical), ice melt will emerge as overland flow with no change in groundwater.
	Wetlands	Increasing inflow to wetlands/peatlands and small lakes which may cause them to expand. Variable change in moisture content of wetland and peatland areas. Changes to water and energy balance. Draining of thermokarst lakes.	As permafrost degrades, infiltration to groundwater will increase and surface runoff will decrease. These changes will increase groundwater storage and flow, yielding greater discharge to wetlands in polar regions. Flushing of preserved contaminants. Secondary thermokarst lakes forming.
	Grasslands	Groundwater will be deep enough to permit little upward connectivity. Groundwater/surface water interactions dominated by infiltration.	As permafrost degrades and surface water is no longer held near the surface, the groundwater table will recede to depths greater than boreal tree species can reach, increasing the occurrence of ecosystems appropriate for grasslands.
River and lake ice	Arctic tundra	River and lake ice will continue to form every winter. May be thinner in rivers that start to maintain continuous flow through the winter. Lake ice may be thinner as winter snowfall increases.	With greater snowpack and insulation from the cold, ice thickness will decrease.
	Boreal plains	Ice thickness in ponds likely to decrease in total thickness. Ice cover in rivers and streams likely to become thinner with more frequent occurrences of midwinter open water.	Ice thickness likely to decrease due to longer periods of groundwater input, greater total input, and increased snow depths.
	Mountains	Ice on mountain lakes likely to decrease in thickness and duration. Ice changes on streams dependent upon terrain and source.	Air temperatures and snow depth will decrease end of winter lake ice thickness, but terrain controls on mountain streams ice will likely not change. Winter icings (or aufeis) could increase markedly if sources of groundwater flow later into the winter.
	Shields	Ice on lakes in shield region is likely to decrease in thickness and duration. Ice on rivers and streams will decrease in	Thicker snowpack, milder temperatures, and greater input of potentially warmer groundwater will contribute to decreased ice.

Table 2. (continued)

Process	Hydrophysiographical Region	Projected Changes	Specific Effects/Comments
		thickness with more frequent midwinter open water.	
	Ice sheets, ice caps, and glaciers	NA	
	Wetlands	Ice on lakes in wetlands is likely to decrease in thickness and duration. Ice on rivers and streams will decrease in thickness with more frequent midwinter open water.	Thicker snowpack, milder temperatures, and greater input of potentially warmer groundwater will contribute to decreased ice.
	Grasslands	Ice on lakes in grasslands is likely to decrease in thickness and duration. Ice on rivers and streams will decrease in thickness with more frequent midwinter open water.	Thicker snowpack, milder temperatures, and greater input of potentially warmer groundwater will contribute to decreased ice.

Vihma *et al.*, 2016]. Based on suite of nine general circulation model (GCM) simulations from CMIP3 over the terrestrial pan-Arctic, Rawlins *et al.* [2010] showed that over the 100 year period from 1950 to 2049, annual evapotranspiration trends range from 0.24 mm yr⁻² to as much as 0.92 mm yr⁻², with the multimodel mean trend at 0.65 mm yr⁻². In general, results suggest acceleration in evapotranspiration over the latter half of the present century.

GCMs generally indicate higher water flows in the future [Kattsov *et al.*, 2005, 2007; Holland *et al.*, 2007; Rawlins *et al.*, 2010] (see also detailed discussion about climate models in Lique *et al.* [2016]). Analyses using a macroscale hydrological model with forcing data from GCMs show projected increases in discharge of up to 31% by the 2080s [Arnell, 2005]. Simulations with a global hydrological model indicate consistent increases of 25–50% across forcing data from three GCMs over the most of the pan-Arctic, except in the lower Ob and Hudson Bay drainages where some decreases are projected [van Vliet *et al.*, 2013]. Similar results were obtained with a larger set of GCM forcing data in a study by Koirala *et al.* [2014]. Model consistency is generally higher for increases than decreases in the Arctic. In contrast to historical flows discussed in section 3.1 above, simulations indicate that future flows will be increasingly driven by evaporation increases from retreating ice cover in the Arctic [Bintanja and Selten, 2014] (see also discussion in Vihma *et al.* [2016]). Projections for high and low flows (expressed as 95th and 10th percentiles of daily values) diverge, with increases of 25–50% consistently projected for eastern Siberia and high Arctic North America and decreases of 0–25% mostly projected for western Siberia, lower latitude North American drainage, and for high flows, also Scandinavia [van Vliet *et al.*, 2013]. However, as the climate forcing is the most important control on the hydrological model output [Arnell, 2005], and as climate models do generally not represent permafrost dynamics adequately [Koven *et al.*, 2013; Slater and Lawrence, 2013], further studies would be needed to confirm these results—for example, results of Koirala *et al.* [2014] indicate a greater occurrence of increases in low flows. Furthermore, controls on minimum flows are complex and scale dependent [Rennermalm *et al.*, 2012], so the low flow results are likely not true for basins of all sizes in the described regions. In general, however, a tendency of increasing river flows is likely, which will raise the transport rates of nutrients, sediment, and carbon in Arctic rivers. Older carbon will also be increasingly mobilized [Aiken *et al.*, 2014]. In addition, north-south gradients of flow availability may change, increasing pressure on water resources in southern basins (see further discussion in Instanes *et al.* [2016] and Vihma *et al.* [2016]).

Irrespective of emission scenario, Arctic permafrost area is projected to decline at a semilinear rate until the 2040s, when decline decelerates in lower emission scenarios. In a high-emission scenario, decline continues at approximately the same rate until 2100 [Slater and Lawrence, 2013; see also Lique *et al.*, 2016]. The ongoing permafrost thaw triggers hydrological regime changes in many ways. With a thickening active layer, some areas that currently store water will change into ones that produce runoff, but trends are different for various landscapes. Furthermore, warming can affect soils in opposing directions, depending on the hydrologic gradient: additional meltwater may cause soil humidification, but it could also be the initial step toward soil drying due to erosion and streamflow intensification. Wetting is more likely in lowlands and plateaus and drying more likely on steeper slopes, where erosion may also increase. Apart from warming, precipitation also influences permafrost, with high prewinter rainfall and snowfall accelerating soil warming through greater latent heat of freezing and greater snow insulation [Iijima *et al.*, 2010]. The temperature of rain is also important, and as rain becomes more frequent, its higher heat content in comparison with snow will influence the melting of

snow and the active layer thickness. *Walvoord et al.* [2012] showed that diminishing permafrost increases the spatial extent of groundwater discharge in lowlands and decreases the proportion of suprapermafrost (shallow) groundwater contribution to total base flow.

Lake ice cover will decrease, according to a recent modeling study, in both thickness (−10 to −50 cm) and duration (−15 to −50 days) during 2040–2079 compared to 1960–1999 [*Dibike et al.*, 2011]. The largest changes are projected for the Pacific coastal regions of North America, northeastern Canada, eastern Europe, Scandinavia, and northern Russia. The snow depth on lake ice is projected to change by −20 to +10 cm and the amount of white ice (i.e., ice forming from wet snow on top of ice) by −20 to +5 cm. In the high latitudes, white ice may form more easily in the future due to increasing snowfall and thinner ice cover [see discussion of changing lake ice effects in *Instanes et al.*, 2016; *Vihma et al.*, 2016; and *Wrona et al.*, 2016].

A future reduction in thermal gradients along northward flowing and ice-covered Arctic rivers has been suggested to decrease spring flooding because of lessening in the severity of ice jamming [*Prowse et al.*, 2010]. On the other hand, snow water equivalent in spring is projected to increase at very high latitudes, particularly in areas with winter temperatures below −30°C [*Adam et al.*, 2009]. The net result of these two factors (magnitude of spring snow water equivalent and severity of ice jams) remains to be quantified but will vary by river basin according to spatial and temporal variability in future precipitation accumulation and snowmelt regimes around the circumpolar north, including in the headwaters of the large basins located in more southerly latitudes [*Prowse et al.*, 2011]. The effects of changes in river ice breakup regimes on freshwater ecosystems and northern infrastructure are reviewed in *Wrona et al.* [2016] and *Instanes et al.* [2016], respectively.

4.2. Hydrophysiographic Regions

Across the tundra, climate warming will generally deepen the active layer and induce a loss of ground ice. Depending on the ice content, places with ice-rich permafrost will undergo subsidence, which leads to thermokarst and small lake formation that affect surface storage and evaporation [*Smith et al.*, 2005; *Jorgenson et al.*, 2008]. A thickened active layer will accommodate more groundwater at the expense of surface runoff, and the magnitude of spring floods may decrease. Time is an important factor, as landscapes will transition from initial wetting due to subsidence to later drying from more complete permafrost degradation and talik formation, depending on the local hydraulic gradient. Snow depths are generally projected to increase due to increasing winter precipitation [*Callaghan et al.*, 2011; see also *Vihma et al.*, 2016]. The southern margins of Arctic tundra will continue to undergo change to shrubs [*Sturm et al.*, 2001], with attendant effects on snow catch, melt, evaporation, and herbivores' access to food, shelter and migration, and shifts in ecosystems. In addition, overland transportation, e.g., snow and ice roads, may be affected with increasing hindrance from shrub growth and shorter snow cover period.

In the boreal plains, warming will also deepen the active layers in permafrost regions, and lateral attrition will take place along the edges of permafrost bodies. In some areas, drying of lakes and wetlands will lead to a browning of forests, a decline in tree growth, and a reduction in latent heat flux, although effects of scale and hydraulic gradient will also influence this process. There is evidence of recent browning of the boreal forests in Central Alaska, which may be related to increased drought stress, though the infestation of insects can also be a factor [*Parent and Verbyla*, 2010; *Verbyla*, 2011]. On the other hand, deepening and moistening of the active layer may also create perennially waterlogged conditions that suppress forest development, something that has been observed in the Lena River delta [*Iijima et al.*, 2014]. Increasing connectivity between surface water and groundwater may support wetland development in areas of groundwater upwelling. Additional implications of increased connectivity include impacts on erosion, sediment, and nutrient release, processes that in turn affect downstream deltas [*Rachold et al.*, 1996, 2000; *Syvitski et al.*, 2005; *Frey and McClelland*, 2009; *Rowland et al.*, 2010] and form new riparian hydrological regimes. Snowmelt timing change will affect plant growth patterns and through them also evapotranspiration and soil moisture. Permafrost degradation may regionally be accelerated by anthropogenic reductions or removal of insulating peat layers. In the southern parts of the boreal plains, human water demand is likely to increase both for natural resource extraction, such as tar sands development and fracking, and agricultural and industrial use [see *Instanes et al.*, 2016]. Fires are likely to increase in frequency [*Stocks et al.*, 1998; *Flannigan et al.*, 2005; *Hu et al.*, 2015]. Possible hydrological effects, either from fires or managed clear-cutting, include higher snow accumulation and subsequent higher volumes of spring meltwater, but such effects are not consistently observed [*Ellis et al.*, 2013; *Semenova et al.*, 2015].

Bedrock terrain can shed snowmeltwater effectively, and the reduced summer rainfall will leave the shields with diminished moisture available for evaporation. However, climate warming can lead to higher evapotranspiration and eutrophication. Depending on the proportion of exposed bedrock and the coverage of boreal vegetation, the rates are spatially variable. With an increase in aridity in parts of the sub-Arctic, a loss of connectivity between the lakes, wetlands, and rivers will be exacerbated in some regions. This has significant implications for the seasonal reduction or loss of flow, with attendant effects on water quality and ecology. Zones of groundwater recharge and discharge, however, depend mostly on the fracture networks in the bedrock and will not change.

In the mountains, the existence of permafrost in subalpine zones reduces storage and enhances runoff generation, but the shrinkage of alpine permafrost can alter this situation. Climate warming and a shift in the rain-snow regime will affect the streamflow regime. A reduction of permafrost in mountain basins will affect components of the water balance of river basins. It has long been recognized that small basins with a low percentage of permafrost produce lower rainfall peaks but higher recession flows than basins underlain by larger amounts of permafrost [Slaughter *et al.*, 1983]. In a changing climate, amplified warming with elevation [Mountain Research Initiative EDW Working Group, 2015] may reinforce changes in Arctic mountain environments.

For grasslands in Canada, one study of a small watershed showed no projected change in annual evapotranspiration and water yield from 1971–2000 to 2041–2070 but a decrease in storage [Zhang *et al.*, 2011]. However, seasonal changes were not consistent, with uncertainties stemming from the choice of climate model. In general, future warming and continuing growth of water abstractions will act to exacerbate water shortage in the grasslands used for agriculture.

Glaciers in the pan-Arctic will mostly continue to thin and retreat, even without additional warming, as they are generally not in balance with the current climate. Regional differences and uncertainties are large, with mean glacier losses between $66 \pm 51\%$ and $11 \pm 28\%$ required to return to equilibrium [Mernild *et al.*, 2014]. A continued lengthening of the ice melt period and possibly warmer summers will raise the elevation of the equilibrium line of glaciers and increase the rate of mass loss. For the pan-Arctic, in general, water fluxes from glaciers will therefore first increase over the near term but will decline as the size of glaciers continue to diminish [Arctic Monitoring and Assessment Program, 2012].

In wetlands, a deepening of the active layer may initially lead to increased storage capacity for additional water and to the formation of small lakes. In the long term, however, continued permafrost degradation may lead to drying out of some wetlands, depending on permafrost conditions and hydraulic gradient. Runoff patterns will change erosion rates and possibly give rise to flushing of stored contaminants. The changing water balance of wetlands and peatlands will also influence the energy balance and the hydrochemical characteristics of catchments.

5. Cross-System Effects

Many of the changes outlined in sections 3 and 4 have effects that go beyond terrestrial hydrology to also influence other components of the Arctic freshwater system. In this section, we describe a number of such links, some of which are also schematically highlighted in Figure 4. We stress that Figure 4 is not intended to thoroughly describe these links, which are complex and often bidirectional, but rather to present an overarching view of some points of cross-system interactions. Companion figures that highlight overarching links in a similar way for other components of the AFS are available in Carmack *et al.* [2016], Instanes *et al.* [2016], Vihma *et al.* [2016], and Wrona *et al.* [2016]. Some examples of more detailed and elaborate system diagrams related to freshwater in the Arctic can be found in Francis *et al.* [2009] and Karlsson *et al.* [2011].

5.1. Processes

The broad increase in winter precipitation over the terrestrial Arctic (with regional summer decreases) may have contributed to the overall increase in the aggregate discharge into the Arctic Ocean [Troy *et al.*, 2012]. At the same time, modeling simulations indicate that a small decrease in summer precipitation along with increased potential for evapotranspiration due to the warming climate has resulted in reduced summer soil moisture in various regions of the terrestrial Arctic [Dirmeyer *et al.*, 2013], with cascading effects for structure and function of ecosystems [see Wrona *et al.*, 2016]. The insulating effect of increased snowpack resulting from increased winter precipitation in the high Arctic may result in permafrost warming and increase in



Figure 4. Links between terrestrial hydrology and other components of the Arctic freshwater system. Links are colored depending on whether they are primarily pertaining to permafrost degradation (orange), decrease in snow and ice cover (blue), more surface water flow (purple), or drier or wetter soils (gray). This graphic is neither meant to be comprehensive nor detailed but is intended to illustrate, at a very overarching level, the multifaceted nature of change in the Arctic freshwater system.

active layer thickness, also with effects on ecosystems and resources [see *Instanes et al.*, 2016]. Moreover, the decrease in the proportion of snowfall (versus rainfall) in the low Arctic and the corresponding increase in snowfall in the high Arctic, as well as the decrease in snow cover duration, affect the seasonality of runoff and magnitude of peak flows in Arctic draining rivers, which also has effects on ecosystems, atmospheric feedbacks, and resources.

Evapotranspiration is the largest vertical freshwater flux out of the Arctic drainage basins and therefore the main link from river basins to the atmosphere [see also *Vihma et al.*, 2016]. Through local evapotranspiration-precipitation recycling, it also constitutes a link back to surface waters, a component that may change with both water reservoir construction and irrigation [*van der Ent et al.*, 2010; *Jarsjö et al.*, 2012; see also *Instanes et al.*, 2016] and soil moisture change [*Hinzman et al.*, 2013]. The latter process is also associated with ecosystem changes [*Wrona et al.*, 2016]. Shifts in vegetation type may increase evapotranspiration by 1–13% until the 2050s across the pan-Arctic [*Pearson et al.*, 2013]. Even without vegetation shifts, increases in evapotranspiration through increasing lake areas may contribute to some observed discrepancies between atmospheric and land surface trends [*Karlsson et al.*, 2015].

River flows strongly affect ecosystems [*Wrona et al.*, 2016] and resources [*Instanes et al.*, 2016], in addition to their flux to the Arctic Ocean, which has implications for thermohaline circulation and marine ecosystems

[see Carmack *et al.*, 2016]. Deltas, lakes, and wetlands are partly fed by runoff, which will govern water levels and impact aquatic and terrestrial life. Hydropower is fundamentally dependent on the supply of freshwater through rivers but also gives rise to considerable seasonal shifts in discharge. The variability of flow, as well as high and low flows, affects both ecosystems [Wrona *et al.*, 2016] and resources [Instanes *et al.*, 2016], including the use of liquid freshwater as drinking water for the Arctic population [White *et al.*, 2007; Alessa *et al.*, 2008; Nilsson *et al.*, 2013; Bring *et al.*, 2015b].

Large-scale atmospheric patterns also influence flow. More intense, positive phases of the Pacific Decadal Oscillation have been shown in observations to be a factor in above-average winter precipitation in the Alaska coast range but below average in interior Alaska, Canada, and eastern Siberia, and subsequently, decreased river discharge in northern Canada [Mantua and Hare, 2002; Déry and Wood, 2005].

Changes in permafrost hydrology are coupled to effects on soil moisture, evapotranspiration, flow pathways, and storage, both in groundwater and surface water, and in ground ice. Where permafrost changes are widespread across the landscape, hydrological effects include modified exchanges of moisture with the atmosphere, lake changes that affect ecosystems, changes in stability and flow patterns that affect resources, and biogeochemical fluxes and erosion that affect the ocean.

A reduction in river and lake ice will influence transportation opportunities in cold and remote regions [see Instanes *et al.*, 2016]. Given that ice-covered freshwater bodies comprise a significant portion of the high-latitude, sub-Arctic and tundra landscape, they need to be considered in climate modeling of this region [Prowse *et al.*, 2011; Lique *et al.*, 2016; Vihma *et al.*, 2016].

5.2. Hydrophysiographic Regions

The Arctic tundra has a direct connection to the riverine coastal domain that encircles the Arctic Ocean. Many small to medium size rivers that originate in the tundra flow directly to the ocean, but the flow magnitude of most of them remains unknown. In the high Arctic, ongoing resource development concentrated along the coast [Andreeva, 1998] may give rise to large local effects on hydrology. Over tundra areas, in general, the exchange with the atmosphere through the boundary layer energy balance is strongly affected by snow and the surface albedo of wet or dry tundra.

In boreal and shield regions, hydropower development has also been extensive, and several large rivers in the Canadian Shield have been dammed or their flows altered by diversion [Déry *et al.*, 2005]. The Nelson River yields 94 km^3 of flow each year, a large portion of which used to previously flow through the Churchill River to the Hudson Bay. After 1980, the hydropower projects of the Nelson were surpassed by a reservoir scheme on La Grande River in Québec, which underwent diversion and now discharges $100 \text{ km}^3 \text{ yr}^{-1}$, compared to its original natural flow of $67 \text{ km}^3 \text{ yr}^{-1}$. Other projects, like some in Norway and Northern Ontario that are less massive in scale, similarly affect river flow, with associated effects of water use on the hydrologic regime, water temperature, and ecology.

Mountains exert strong influence on atmospheric circulation and precipitation, both over synoptic and local scales. Many mountain streams are intensely modified by hydropower regulation, which in turn influences downstream ecosystems and seasonal water balances. Other effects on resources include a changing snow cover duration, which will influence recreational activities such as skiing, and also avalanches that pose hazards to transportation and settlements. Furthermore, heavy rainfall can cause landslides that affect traffic and infrastructure in the Norwegian and Cordilleran mountains.

Similar to shield and boreal regions, river systems in grasslands have in many places been subject to extensive modification and diversions [Yang *et al.*, 2004a, 2004b; Schindler and Donahue, 2006; Stuefer *et al.*, 2011], which influence both water exchanges with the atmosphere, final transport to the ocean, and local ecosystems and resources. In Canadian grasslands, paleoclimatic evidence suggests that the relatively moist conditions that prevailed during the twentieth century are unusual [Sauchyn *et al.*, 2002] and that the now heavily modified water systems may become subject to severe water shortages in summer if the prolonged droughts of preindustrial times were to return in combination with elevated pressures from climate change [Schindler and Donahue, 2006].

The mass loss of Arctic glaciers and the GrIS are presently the largest contributors to ocean storage and global sea level rise. In addition, changes in the GrIS outflow have potential ramifications for ocean thermohaline

circulation around Greenland [Rahmstorf *et al.*, 2005, 2015; see also Carmack *et al.*, 2016]. Glaciers are important suppliers of freshwater to mountainous basins, and their shrinkage will require changed planning of hydropower operations in the long run [see Instanes *et al.*, 2016]. Locally, long-term streamflow and flooding increases may be amplified in glacierized basins [Dahlke *et al.*, 2012]. Spreading and thickening of aufeis [Kane, 1981] may be intensified due to increase of underground water storage and release, in turn caused by permafrost thaw.

Arctic wetland systems may act as accumulators of water, where it is increasingly subject to evapotranspiration. They also constitute important water storage reservoirs that regionally are strong controls on runoff formation. Most Arctic wetlands are generally species-rich ecosystems, with marshes and deltas as particularly boundary-crossing regions between terrestrial hydrological regimes and oceans. In this regard, the four largest Arctic rivers have markedly different interfaces with the Arctic Ocean. On the Lena and the Mackenzie, the river flows into the ocean through very large deltas, dominated by the river water. In contrast, the Yenisey and Ob Rivers enter the ocean through marine-dominated estuaries, where tidal flows and surges exert strong influence on local environments.

In estuaries, which are elongated and open for tides, the hydraulic regime is more dynamic than in the deltas. Sedimentation in estuaries occurs under salt-fresh water contact and biogeochemical transformation [Gordeev *et al.*, 1999; Gordeev, 2006; Fedorova *et al.*, 2015]. The marine saltwater intrusion rises higher along the river in estuaries than in deltas and also consists of a more distinct salt-fresh water geochemical barrier. The freshwater plume extending into the ocean from river mouths can therefore be said to be reversed during times in estuaries. In deltas, however, the accumulation of material can be 8–10 times higher than in estuaries. For example, the turbidity in the top of the Lena River delta is 40 mg L^{-1} on average but falls to $3\text{--}5 \text{ mg L}^{-1}$ at the coastal line [Fedorova *et al.*, 2015].

6. Major Knowledge Gaps and Future Research Directions

6.1. Processes

Arctic terrestrial precipitation trends are inherently difficult to detect given snowfall measurement challenges resulting from gauge undercatch of solid precipitation, sparsely distributed observations, low precipitation amounts, and rare long-term records [Serreze and Hurst, 2000; Adam and Lettenmaier, 2003; Yang *et al.*, 2005]. Furthermore, the large natural variability of precipitation implies that a trend may take time to emerge. The compounding effects of elevation on precipitation in topographically complex regions of the Arctic, where the distribution of observing stations is biased toward low elevations and coastal regions, are also a factor [Kane and Stuefer, 2015]. As solid precipitation remains challenging to measure, in situ and remote sensing observations of snow cover on the ground are widely used as a precipitation metric instead of traditional precipitation gauge measurements [Stuefer *et al.*, 2013]. Improving representation of precipitation and snow cover processes specific to the high latitudes, both by better remote sensing technology and more skillful models from a land surface perspective, remains an important research direction that would enhance our ability to understand and model Arctic freshwater [see Lique *et al.*, 2016] and its role in the larger Earth system.

Evapotranspiration measurements in the terrestrial Arctic are mostly from field programs of short duration covering specific regions [Langer *et al.*, 2011; Mueller *et al.*, 2011; see also Vihma *et al.*, 2016]. Because of these data gaps, analysis of evapotranspiration over the terrestrial Arctic often involves information from land surface models and remote sensing data, as well as global reanalysis and climate model outputs [Rawlins *et al.*, 2010]. However, the progress in exploiting remote sensing techniques for measuring high-latitude precipitation and evapotranspiration is very slow and requires increased efforts from the hydrological community [Lettenmaier *et al.*, 2015]. The complexity of modeling evapotranspiration arises from heterogeneity in land surface and a large set of parameters to consider, including soil properties, characteristics of vegetation, and availability of moisture. Studies that examine changes in land surface variables and fluxes [e.g., Destouni *et al.*, 2013; Dirmeyer *et al.*, 2013; Jaramillo and Destouni, 2014; Törnqvist *et al.*, 2014; van der Velde *et al.*, 2014; Bring *et al.*, 2015a], but focus on the terrestrial Arctic, may improve our picture of the land-atmosphere interactions with respect to soil moisture and evapotranspiration. In general, there is a great need for more accurate representation of land surface effects, such as vegetation, lakes, and human interventions, in climate models [Harding *et al.*, 2014; Bring *et al.*, 2015a].

Knowledge of runoff variability in the Arctic is limited by river observations, which cover about 70% of the total contributing area [Lammers *et al.*, 2001; Shiklomanov *et al.*, 2002, 2006; Bring and Destouni, 2013], and whose availability and reliability are even more lacking when considering water chemistry and sediment [Holmes *et al.*, 2000, 2012; Prowse *et al.*, 2005; McClelland *et al.*, 2008, 2015; Bring and Destouni, 2009]. Notwithstanding these limitations, our understanding of how discharge changes in response to land surface and permafrost changes is not complete. Observed divergence in precipitation and discharge trends across basins [e.g., Berezovskaya *et al.*, 2004; Bring and Destouni, 2011] indicate that changes in surface and subsurface water interactions, and their feedback on water balance and evapotranspiration [Bosson *et al.*, 2013], need to be better understood and quantified, although in some cases such discrepancies are due to data uncertainty. Climate model agreement on runoff changes is also particularly limited for Arctic basins [Bring and Destouni, 2011; Törnqvist *et al.*, 2014]. Along the coast, information shortages combine to yield great uncertainty both in surface flows, which are poorly monitored for many smaller rivers, and the interaction between groundwater, surface water, permafrost, and seawater [Bobba *et al.*, 2012; see also Carmack *et al.*, 2016]. To address these knowledge gaps in river fluxes, some proposed strategies to improve monitoring may help [e.g., Karlsson *et al.*, 2011; Azcárate *et al.*, 2013; McClelland *et al.*, 2015], as will a continuation of the Sustaining Arctic Observation Networks process. The potential of remote sensing methods, as a complement to ground observations, will likely increase with new instruments (e.g., the NASA Surface Water and Ocean Topography mission and the EU Sentinel missions), and some promising results so far [e.g., Smith and Pavelsky, 2008; Biancamaria *et al.*, 2011; Gleason and Smith, 2014] indicate that the Arctic hydrology community should investigate these opportunities further. Another remaining knowledge gap is how to ensure efficient long-term monitoring of Arctic rivers under changing climate conditions, something that should be addressed by combining climate projections and network optimization methods. Furthermore, updates are now required to the rating curves for major Arctic gauges, without which a much desired coordinated monitoring of water temperature, chemistry, and suspended matter will be hampered [McClelland *et al.*, 2015]. Hydrology-ecology interactions should be investigated through field monitoring and new methods using isotopes and remote sensing. Multidisciplinary investigations at the drainage basin scale should be encouraged as they have the potential to increase understanding of interaction between hydrology and atmosphere, oceans, ecology, and resources [see also Wrona *et al.*, 2016].

Main knowledge gaps in permafrost hydrology include the understanding of underground and surface water interaction (for example, inflow and outflow from riverbed taliks), the mechanisms of infiltration and flow formation in frozen and thawing ground in different landscapes, evaporation processes, atmosphere-land surface-subsurface energy interaction, and the patterns of precipitation distribution at catchment scales (including snow redistribution and intensive rain events in mountainous conditions). Scale discrepancies between field studies and catchments, and between hydrological models of various scales and climate data, are a challenge. Runoff formation governed by active layer thaw-freeze processes varies spatially at slope scale, and modeling and parameter estimation, even for large basins, require consideration of this. Advances in modeling these processes, ideally in collaboration with the soil carbon community, would be a key step toward understanding how wetting and drying of permafrost landscapes will influence soil carbon release to the atmosphere during the present century [Schoor *et al.*, 2015]. Observational data uncertainty, especially for extreme conditions, remains an obstacle to advances in understanding. However, existing catchment study data are sometimes underutilized. Additional parameters for small and large models can be assessed based on field measurements and improved equipment. To capitalize on these underexploited opportunities, integrating data and studies across the Arctic should be a priority.

6.2. Hydrophysiographic Regions

An overarching and pressing research question across all regions of the Arctic is how to separate anthropogenic from natural disturbances. Human impacts are not easily quantified as the type and intensity of activities are difficult to predict, and their effects on hydrology are to a degree uncertain, particularly on a larger systems level. The potential for new methods that distinguish change drivers, previously applied globally to separate landscape and atmospheric components [Jaramillo and Destouni, 2014], should be investigated also for the Arctic region.

Anthropogenic activities are particularly likely to increase in the coastal tundra zone [Andreeva, 1998]. However, the land-ocean boundary commonly falls in between central monitoring and observation strategies that prioritize information goals based on either a terrestrial or a marine perspective [Strandmark *et al.*, 2015]. An explicit

focus on the hydrological interactions between land and sea across all of the coastal interface, and not just through rivers, would help overbridge disciplinary boundaries and expand knowledge on a relatively understudied part of the Arctic freshwater system. Also, our understanding of marsh hydrology, including hydrodynamics and hydroecology of the Arctic coastal zone, is limited [Overduin *et al.*, 2014], as well as our knowledge of submarine groundwater discharge and the forming of salt-fresh aquifers in coastal zones [Bobba *et al.*, 2012]. Groundwater models of varying complexity [e.g., Danielescu *et al.*, 2009; Mazi *et al.*, 2013], potentially coupled to hydrological and oceanographic models, should be applied to Arctic conditions to increase understanding of the coastal zone.

Large parts of the boreal zone are underlain by discontinuous permafrost, and these areas are also prone to development in the near future. The anthropogenic impact on the hydrological cycle from forest clearing and fires is not fully understood. More information is needed on the present status of groundwater circulation and storage in discontinuous permafrost terrain in order to understand impacts on heat and water fluxes and reasonably project future conditions. Emerging and improved remote sensing techniques will enhance the opportunity for surveillance of surface wetness on a regional scale [e.g., Leconte *et al.*, 2008]. Future studies of mountain hydrology are also needed to determine if those areas will continue to provide as large a share of river flow to the Arctic Ocean as today.

Projections of future precipitation changes constitute a key hydrological uncertainty in grasslands [Zhang *et al.*, 2011]. In addition, the previously noted uncertainty in future drought patterns [Schindler and Donahue, 2006] will strongly influence grassland hydrology. To improve knowledge and improve adaptation efforts, improved downscaling of drought-related variables, as well as further exploration of teleconnection patterns that govern droughts [Schindler and Donahue, 2006], is necessary. In grasslands, the extensive human modifications of the water cycle should also be considered in both model projections of future changes [Bring *et al.*, 2015a] and in the evaluation of past changes [Jaramillo and Destouni, 2015].

In terms of glaciers and ice sheets, dominant unknowns and uncertainties in modeling and understanding of ice sheet mass balance and runoff response are percolation, retention, and freezing of melt and rain water by snow [e.g., Box, 2013; Vernon *et al.*, 2013]. Extensive surface melt has been recently documented over the GrIS, unprecedented in the observational record (NOAA Arctic report card 2012 available at www.arctic.noaa.gov/report-card) [Nghiem *et al.*, 2012; Hanna *et al.*, 2014]. Meltwater can be trapped within the firn, and so the magnitude and delay of runoff remain unknown. Conventional thought has been that practically all air space in the snow must be filled before meltwater can exit the ice sheets as runoff. Yet recent snow and ice core measurements provide the first evidence of the formation of impermeable ice layers, enabling an abrupt increase in ice sheet runoff [Vernon *et al.*, 2013]. Other recent results indicate that ice sheet outflow may be overestimated in climate models due to omission of subglacial processes [Smith *et al.*, 2015]. Development of percolation/retention schemes and evaluation of nonlinearities in ice sheet meltwater runoff should therefore lead to more accurate projections of GrIS runoff.

To increase our understanding of wetland dynamics, there is a need for improved riverbed morphological parametrization for hydraulic models, especially for large Arctic rivers. Such data would also improve the ability to model river fluxes, in general, and not just wetlands. In particular, there is a need to monitor the seasonal and interannual changes in wetlands, as near-surface moisture conditions strongly affect evaporation and can influence the development of precipitation and even summer weather. Evapotranspiration measurements are critical to understand water fluxes, and organization of such measurements across different wetland and peatland types is required. There is also a need for method improvements, including models, remote sensing, and isotopic techniques [Tetzlaff *et al.*, 2015].

7. Conclusion

Terrestrial freshwater pathways have a central role in water, material (ions), and energy exchanges between components of the Arctic freshwater system. They connect the atmosphere and ocean over large distances, along which ecosystems and resources are affected by varying geographical characteristics of the terrestrial freshwater systems.

Across the terrestrial Arctic, snow cover is generally decreasing, but patterns are complex in terms of duration and extent, and the snow water equivalent. Evapotranspiration is coupled to landscape and permafrost changes in different ways across the Arctic, with wetting and drying occurring in succession, depending on the

permafrost conditions. However, some regions may undergo systematic shifts in one direction or the other. Annual river flux patterns show a long-term increase at the pan-Arctic scale, and patterns of higher discharge also seem to apply more universally across regions in recent decades. Reports of permafrost warming are numerous, and an expected effect of increasing river base flows has been reported for several regions. River and lake ice cover have decreased in duration, with the most rapid reductions observed from the late twentieth century.

Water flux and storage change in complex ways across Arctic regions. Permafrost degradation will occur both in the tundra and boreal zone, with complete permafrost degradation occurring along the margins of the permafrost zone in the taiga. Evapotranspiration and precipitation are likely to increase in all regions, although summer precipitation may decrease in the southern parts of the Arctic. River flows will likely increase on average across all regions, but high and low flows may decrease in parts of the Arctic boreal plains, shields, and grasslands regions, mostly where local decreases in average flows are also projected.

To improve our understanding of changes to the freshwater system in the Arctic, a number of knowledge gaps are of particular importance. We are still lacking a comprehensive picture of evapotranspiration across Arctic regions and how it will interact with climate and landscape changes. These changes are coupled to precipitation, river flows, and groundwater flows, which underlines that it remains a challenge to estimate full water budgets for most Arctic river basins. For coastal regions, the interaction of flow along various poorly constrained pathways implies that these regions are essentially uncharted. As some parts of the Arctic increasingly become subject to settlement and natural resource extraction, it will become more important to separate anthropogenic effects from underlying environmental changes.

To address these challenges, future research should include more widespread use of multimodel ensembles to enable better separation of model-internal variability from change. Novel approaches to overcome data limitations will be needed, as they are not likely to diminish appreciably without large-scale funding increases. Concerted efforts at studying water, solute, and energy fluxes on various scales will remain important to detect, understand, and project water system changes. Despite a growing potential of remote sensing, it remains critical to at least maintain the current ground-based capacity to observe Arctic terrestrial hydrology.

Acknowledgments

The Arctic Freshwater Synthesis has been sponsored by the World Climate Research Programme's Climate and the Cryosphere project (WCRP-CriC), the International Arctic Science Committee (IASC), and the Arctic Monitoring and Assessment Program (AMAP). Additional support has been provided by the Norwegian Ministries of Environment and of Foreign Affairs, the Swedish Secretariat for Environmental Earth System Sciences (SSEESS), and the Swedish Polar Research Secretariat. We gratefully acknowledge the project coordination and meeting support of Jenny Baeseman and Gwenaëlle Hamon at the CliC International Project Office and thank Brandi Newton and Hayley Linton for providing data analysis for Figure 2. A. Bring acknowledges support from the Swedish Research Council VR (project 2013-7448). O. Semanova acknowledges support from the Russian Foundation for Basic Research (project 15-35-21146). All data for this paper are cited and referred to in the reference list. The data used to produce Figure 2 are available from the ERA-Interim data sets [Dee *et al.*, 2011] (<http://apps.ecmwf.int/datasets/data/interim-full-daily>), and the data to produce Figure 3 are available from the Global Runoff Data Centre (<http://www.bafg.de>) and the R-ArcticNET database (<http://www.r-arcticnet.sr.unh.edu>).

References

- Aagaard, K., and E. C. Carmack (1989), The role of sea ice and other fresh water in the Arctic circulation, *J. Geophys. Res.*, *94*, 14,485–14,498, doi:10.1029/JC094iC10p14485.
- Adam, J. C., and D. P. Lettenmaier (2003), Adjustment of global gridded precipitation for systematic bias, *J. Geophys. Res.*, *108*(D9), 4257, doi:10.1029/2002JD002499.
- Adam, J. C., E. A. Clark, D. P. Lettenmaier, and E. F. Wood (2006), Correction of global precipitation products for orographic effects, *J. Clim.*, *19*(1), 15–38.
- Adam, J. C., A. F. Hamlet, and D. P. Lettenmaier (2009), Implications of global climate change for snowmelt hydrology in the twenty-first century, *Hydrol. Processes*, *23*(7), 962–972.
- Aiken, G. R., R. G. M. Spencer, R. G. Striegel, P. F. Schuster, and P. A. Raymond (2014), Influences of glacier melt and permafrost thaw on the age of dissolved organic carbon in the Yukon River basin, *Global Biogeochem. Cycles*, *28*, 525–537, doi:10.1002/2013GB004764.
- Alessa, L., A. Kliskey, R. Lammers, C. Arp, D. White, L. Hinzman, and R. Busey (2008), The Arctic water resource vulnerability index: An integrated assessment tool for community resilience and vulnerability with respect to freshwater, *Environ. Manage.*, *42*(3), 523–541, doi:10.1007/s00267-008-9152-0.
- Arctic Monitoring and Assessment Program (2012), *Arctic Climate Issues 2011: Changes in Arctic Snow, Water, Ice and Permafrost. SWIPA 2011 Overview Report*, Arctic Monitoring and Assessment Programme (AMAP), Oslo.
- Ananicheva, M. D., and G. Kapustin (2010), Change of glacier state in mountain regions of the Russian Sub-Arctic: Assessment by satellite imagery and aerial photography in the USSR Glacier Inventory-Russian contribution into the IPY. 2007–2008, *Cryosphere*, *3*, 19–26.
- Ananicheva, M. D., G. A. Kapustin, and M. M. Koreysha (2006), Glacier changes in Suntar-Khayata mountains and Chersky Range from the Glacier Inventory of the USSR and satellite images 2001–2003, *Data Glaciol. Stud. Moscow*, *101*, 163–169.
- Anderson, R. K., G. H. Miller, J. P. Briner, N. A. Lifton, and S. B. DeVogel (2008), A millennial perspective on Arctic warming from ¹⁴C in quartz and plants emerging from beneath ice caps, *Geophys. Res. Lett.*, *35*, L01502, doi:10.1029/2007GL032057.
- Andreeva, E. N. (1998), The Russian Arctic coastal zone management problems: Past lessons and new realities, *Ocean Coastal Manage.*, *41*(2-3), 237–256, doi:10.1016/S0964-5691(98)00067-2.
- Andresen, C. G., and V. L. Loughheed (2015), Disappearing Arctic tundra ponds: Fine-scale analysis of surface hydrology in drained thaw lake basins over a 65 year period (1948–2013), *J. Geophys. Res. Biogeosci.*, *120*, 466–479, doi:10.1002/2014JG002778.
- Armstrong, R. N., J. W. Pomeroy, and L. W. Martz (2015), Variability in evaporation across the Canadian Prairie region during drought and non-drought periods, *J. Hydrol.*, *521*, 182–195.
- Arnell, N. W. (2005), Implications of climate change for freshwater inflows to the Arctic Ocean, *J. Geophys. Res.*, *110*, D07105, doi:10.1029/2004JD005348.
- Azcárate, J., B. Balfors, A. Bring, and G. Destouni (2013), Strategic environmental assessment and monitoring: Arctic key gaps and bridging pathways, *Environ. Res. Lett.*, *8*(4), 044033, doi:10.1088/1748-9326/8/4/044033.

- Barrand, N. E., and M. J. Sharp (2010), Sustained rapid shrinkage of Yukon glaciers since the 1957–1958 International Geophysical Year, *Geophys. Res. Lett.*, *37*, L07501, doi:10.1029/2009GL042030.
- Beltaos, S., and T. Prowse (2009), River-ice hydrology in a shrinking cryosphere, *Hydrol. Processes*, *23*(1), 122–144.
- Bennett, K. E., and T. D. Prowse (2010), Northern Hemisphere geography of ice-covered rivers, *Hydrol. Processes*, *24*(2), 235–240.
- Bense, V. F., G. Ferguson, and H. Kooi (2009), Evolution of shallow groundwater flow systems in areas of degrading permafrost, *Geophys. Res. Lett.*, *36*, L22401, doi:10.1029/2009GL039225.
- Bense, V. F., H. Kooi, G. Ferguson, and T. Read (2012), Permafrost degradation as a control on hydrogeological regime shifts in a warming climate, *J. Geophys. Res.*, *117*, F03036, doi:10.1029/2011JF002143.
- Benson, B. J., J. J. Magnuson, O. P. Jensen, V. M. Card, G. Hodgkins, J. Korhonen, D. M. Livingstone, K. M. Stewart, G. A. Weyhenmeyer, and N. G. Granin (2011), Extreme events, trends, and variability in Northern Hemisphere lake-ice phenology (1855–2005), *Clim. Change*, *112*(2), 299–323, doi:10.1007/s10584-011-0212-8.
- Berezovskaya, S., D. Yang, and D. L. Kane (2004), Compatibility analysis of precipitation and runoff trends over the large Siberian watersheds, *Geophys. Res. Lett.*, *31*, L21502, doi:10.1029/2004GL021277.
- Berezovskaya, S., D. Yang, and L. Hinzman (2005), Long-term annual water balance analysis of the Lena River, *Global Planet. Change*, *48*, 84–95, doi:10.1016/j.gloplacha.2004.12.006.
- Biancamaria, S., M. Durand, K. M. Andreadis, P. D. Bates, A. Boone, N. M. Mognard, E. Rodriguez, D. E. Alsdorf, D. P. Lettenmaier, and E. A. Clark (2011), Assimilation of virtual wide swath altimetry to improve Arctic river modeling, *Remote Sens. Environ.*, *115*(2), 373–381, doi:10.1016/j.rse.2010.09.008.
- Bintanja, R., and F. M. Selten (2014), Future increases in Arctic precipitation linked to local evaporation and sea-ice retreat, *Nature*, *509*(7501), 479–482, doi:10.1038/nature13259.
- Bobba, A. G., P. A. Chambers, and F. J. Wrona (2012), Submarine groundwater discharge (SGWD): An unseen yet potentially important coastal phenomenon in Canada, *Nat. Hazards*, *60*(3), 991–1012, doi:10.1007/s11069-011-9884-7.
- Bolch, T., B. Menounos, and R. Wheate (2010), Landsat-based inventory of glaciers in western Canada, 1985–2005, *Remote Sens. Environ.*, *114*(1), 127–137.
- Bosson, E., J.-O. Selroos, M. Stigsson, L.-G. Gustafsson, and G. Destouni (2013), Exchange and pathways of deep and shallow groundwater in different climate and permafrost conditions using the Forsmark site, Sweden, as an example catchment, *Hydrogeol. J.*, *21*(1), 225–237.
- Bowling, L. C., and D. P. Lettenmaier (2010), Modeling the effects of lakes and wetlands on the water balance of Arctic environments, *J. Hydrometeorol.*, *11*(2), 276–295, doi:10.1175/2009JHM1084.1.
- Bowling, L. C., D. P. Lettenmaier, and B. V. Matheussen (2000), Hydroclimatology of the Arctic drainage basin, in *The Freshwater Budget of the Arctic Ocean*, edited by E. L. Lewis et al., pp. 57–90, Springer, Dordrecht, Netherlands.
- Box, J. E. (2013), Greenland ice sheet mass balance reconstruction. Part II: Surface mass balance (1840–2010), *J. Clim.*, *26*(18), 6974–6989.
- Box, J. E., and W. Colgan (2013), Greenland ice sheet mass balance reconstruction. Part III: Marine ice loss and total mass balance (1840–2010), *J. Clim.*, *26*(18), 6990–7002.
- Bring, A., and G. Destouni (2009), Hydrological and hydrochemical observation status in the pan-Arctic drainage basin, *Polar Res.*, *28*, 327–338, doi:10.1111/j.1751-8369.2009.00126.x.
- Bring, A., and G. Destouni (2011), Relevance of hydro-climatic change projection and monitoring for assessment of water cycle changes in the Arctic, *Ambio*, *40*(4), 361–369, doi:10.1007/s13280-010-0109-1.
- Bring, A., and G. Destouni (2013), Hydro-climatic changes and their monitoring in the Arctic: Observation-model comparisons and prioritization options for monitoring development, *J. Hydrol.*, *492*, 273–280, doi:10.1016/j.jhydrol.2013.04.003.
- Bring, A., and G. Destouni (2014), Arctic climate and water change: Model and observation relevance for assessment and adaptation, *Surv. Geophys.*, *35*, 853–877, doi:10.1007/s10712-013-9267-6.
- Bring, A., S. M. Asokan, F. Jaramillo, J. Jarsjö, L. Levi, J. Pietron, C. Prieto, P. Rogberg, and G. Destouni (2015a), Implications of freshwater flux data from the CMIP5 multimodel output across a set of Northern Hemisphere drainage basins, *Earths Future*, *3*(6), 206–217, doi:10.1002/2014EF000296.
- Bring, A., J. Jarsjö, and G. Destouni (2015b), Water information and water security in the Arctic, in *The New Arctic*, edited by B. Evengård, J. Nymand Larsen, and Ø. Paasche, Springer, Cham, Switzerland.
- Brooks, R. N., T. D. Prowse, and I. J. O'Connell (2013), Quantifying Northern Hemisphere freshwater ice, *Geophys. Res. Lett.*, *40*, 1128–1131, doi:10.1002/grl.50238.
- Brown, R. D., and P. W. Mote (2009), The response of Northern Hemisphere snow cover to a changing climate, *J. Clim.*, *22*(8), 2124–2145, doi:10.1175/2008JCLI2665.1.
- Brown, R. D., and D. A. Robinson (2011), Northern Hemisphere spring snow cover variability and change over 1922–2010 including an assessment of uncertainty, *Cryosphere*, *5*(1), 219–229.
- Brutel-Vuilmet, C., M. Ménégoz, and G. Krinner (2013), An analysis of present and future seasonal Northern Hemisphere land snow cover simulated by CMIP5 coupled climate models, *Cryosphere*, *7*, 67–80.
- Brutsaert, W., and T. Hiyama (2012), The determination of permafrost thawing trends from long-term streamflow measurements with an application in eastern Siberia, *J. Geophys. Res.*, *117*, D22110, doi:10.1029/2012JD018344.
- Callaghan, T. V., et al. (2011), The changing face of Arctic snow cover: A synthesis of observed and projected changes, *Ambio*, *40*(1), 17–31, doi:10.1007/s13280-011-0212-y.
- Carey, S. K., and M. Woo (2001), Slope runoff processes and flow generation in a subarctic, subalpine catchment, *J. Hydrol.*, *253*(1), 110–129.
- Carmack, E. C., et al. (2016), Fresh water and its role in the Arctic Marine System: Sources, disposition, storage, export, and physical and biogeochemical consequences in the Arctic and global oceans, *J. Geophys. Res. Biogeosci.*, *121*, doi:10.1002/2015JG003140.
- Church, J. A., et al. (2013), Sea level change, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by T. F. Stocker et al., pp. 1137–1216, Cambridge Univ. Press, Cambridge, U. K.
- Cogley, J. G. (2012), The future of the world's glaciers, in *Future World's Climate*, 2nd ed., pp. 197–222, Elsevier, Amsterdam.
- Connon, R. F., W. L. Quinton, J. R. Craig, and M. Hayashi (2014), Changing hydrologic connectivity due to permafrost thaw in the lower Liard River valley, NWT, Canada, *Hydrol. Processes*, *28*(14), 4163–4178, doi:10.1002/hyp.10206.
- Dahlke, H. E., S. W. Lyon, J. R. Stedinger, G. Rosqvist, and P. Jansson (2012), Contrasting trends in floods for two sub-arctic catchments in northern Sweden—Does glacier presence matter?, *Hydrol. Earth Syst. Sci.*, *16*(7), 2123–2141.
- Danby, R. K., and D. S. Hik (2007), Evidence of recent treeline dynamics in southwest Yukon from aerial photographs, *Arctic*, *60*, 411–420.
- Danielescu, S., K. T. B. MacQuarrie, and R. N. Faux (2009), The integration of thermal infrared imaging, discharge measurements and numerical simulation to quantify the relative contributions of freshwater inflows to small estuaries in Atlantic Canada, *Hydrol. Processes*, *23*(20), 2847–2859, doi:10.1002/hyp.7383.

- Dankers, R. (2003), Sub-arctic hydrology and climate change: A case study of the Tana River Basin in Northern Fennoscandia, Thesis, Utrecht Univ., Utrecht, Netherlands.
- Dee, D. P., et al. (2011), The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Q. J. R. Meteorol. Soc.*, *137*(656), 553–597.
- De Rham, L. P., T. D. Prowse, S. Beltaos, and M. P. Lacroix (2008), Assessment of annual high-water events for the Mackenzie River basin, Canada, *Hydrol. Processes*, *22*(18), 3864–3880.
- Déry, S. J., and E. F. Wood (2005), Decreasing river discharge in northern Canada, *Geophys. Res. Lett.*, *32*, L10401, doi:10.1029/2005GL022845.
- Déry, S. J., M. Stieglitz, E. C. McKenna, and E. F. Wood (2005), Characteristics and trends of river discharge into Hudson, James, and Ungava Bays, 1964–2000, *J. Clim.*, *18*(14), 2540–2557.
- Déry, S. J., M. A. Hernández-Henríquez, J. E. Burford, and E. F. Wood (2009), Observational evidence of an intensifying hydrological cycle in northern Canada, *Geophys. Res. Lett.*, *36*, L13402, doi:10.1029/2009GL038852.
- Déry, S. J., T. J. Mlynowski, M. A. Hernández-Henríquez, and F. Straneo (2011), Interannual variability and interdecadal trends in Hudson Bay streamflow, *J. Mar. Syst.*, *88*(3), 341–351.
- Deser, C., R. Tomas, M. Alexander, and D. Lawrence (2010), The seasonal atmospheric response to projected Arctic sea ice loss in the late twenty-first century, *J. Clim.*, *23*(2), 333–351.
- Destouni, G., F. Jaramillo, and C. Prieto (2013), Hydroclimatic shifts driven by human water use for food and energy production, *Nat. Clim. Change*, *3*, 213–217, doi:10.1038/nclimate1719.
- Dibike, Y., T. Prowse, T. Saloranta, and R. Ahmed (2011), Response of Northern Hemisphere lake-ice cover and lake-water thermal structure patterns to a changing climate, *Hydrol. Processes*, *25*(19), 2942–2953.
- Dirmeyer, P. A., Y. Jin, B. Singh, and X. Yan (2013), Trends in land–atmosphere interactions from CMIP5 simulations, *J. Hydrometeorol.*, *14*(3), 829–849.
- Dyrgerov, M. B., A. Bring, and G. Destouni (2010), Integrated assessment of changes in freshwater inflow to the Arctic Ocean, *J. Geophys. Res.*, *115*, D12116, doi:10.1029/2009JD013060.
- Edlund, S. A., and B. A. Alt (1989), Regional congruence of vegetation and summer climate patterns in the Queen Elizabeth Islands, Northwest Territories, Canada, *Arctic*, *42*, 3–23.
- Ehsanzadeh, E., and K. Adamowski (2010), Trends in timing of low stream flows in Canada: Impact of autocorrelation and long-term persistence, *Hydrol. Processes*, *24*(8), 970–980, doi:10.1002/hyp.7533.
- Ellis, C. R., J. W. Pomeroy, and T. E. Link (2013), Modeling increases in snowmelt yield and desynchronization resulting from forest gap-thinning treatments in a northern mountain headwater basin, *Water Resour. Res.*, *49*, 936–949, doi:10.1002/wrcr.20089.
- Fang, X., and J. W. Pomeroy (2008), Drought impacts on Canadian prairie wetland snow hydrology, *Hydrol. Processes*, *22*(15), 2858–2873, doi:10.1002/hyp.7074.
- Fedorova, I., et al. (2015), Lena Delta hydrology and geochemistry: Long-term hydrological data and recent field observations, *Biogeosciences*, *12*(2), 345–363.
- Ferone, J. M., and K. J. Devito (2004), Shallow groundwater–surface water interactions in pond–peatland complexes along a Boreal Plains topographic gradient, *J. Hydrol.*, *292*(1–4), 75–95, doi:10.1016/j.jhydrol.2003.12.032.
- Flannigan, M. D., K. A. Logan, B. D. Amiro, W. R. Skinner, and B. J. Stocks (2005), Future area burned in Canada, *Clim. Change*, *72*(1–2), 1–16.
- Frampton, A., S. Painter, S. W. Lyon, and G. Destouni (2011), Non-isothermal, three-phase simulations of near-surface flows in a model permafrost system under seasonal variability and climate change, *J. Hydrol.*, *403*(3–4), 352–359, doi:10.1016/j.jhydrol.2011.04.010.
- Frampton, A., S. L. Painter, and G. Destouni (2013), Permafrost degradation and subsurface-flow changes caused by surface warming trends, *Hydrogeol. J.*, *21*(1), 271–280, doi:10.1007/s10040-012-0938-z.
- Francis, J. A., D. M. White, J. J. Cassano, W. J. Gutowski, L. D. Hinzman, M. M. Holland, M. A. Steele, and C. J. Vörösmarty (2009), An arctic hydrologic system in transition: Feedbacks and impacts on terrestrial, marine, and human life, *J. Geophys. Res.*, *114*, G04019, doi:10.1029/2008JG000902.
- Frey, K. E., and J. W. McClelland (2009), Impacts of permafrost degradation on arctic river biogeochemistry, *Hydrol. Processes*, *23*(1), 169–182.
- Gleason, C. J., and L. C. Smith (2014), Toward global mapping of river discharge using satellite images and at-many-stations hydraulic geometry, *Proc. Natl. Acad. Sci. U. S. A.*, *111*(13), 4788–4791, doi:10.1073/pnas.1317606111.
- Gordeev, V. V. (2006), Fluvial sediment flux to the Arctic Ocean, *Geomorphology*, *80*, 94–104.
- Gordeev, V. V., I. S. Zektser, R. G. Djmalov, A. V. Zhulidov, and V. A. Bryzgalov (1999), Score removal of nutrients of river runoff to the marginal seas of the Russian Arctic, *Water Resour.*, *26*, 206–211.
- Goulding, H. L., T. D. Prowse, and S. Beltaos (2009), Spatial and temporal patterns of break-up and ice-jam flooding in the Mackenzie Delta, NWT, *Hydrol. Processes*, *23*(18), 2654–2670.
- Haine, T. W. N., et al. (2015), Arctic freshwater export: Status, mechanisms, and prospects, *Global Planet. Change*, *125*, 13–35, doi:10.1016/j.gloplacha.2014.11.013.
- Hanna, E., P. Huybrechts, K. Steffen, J. Cappelien, R. Huff, C. Shuman, T. Irvine-Fynn, S. Wise, and M. Griffiths (2008), Increased runoff from melt from the Greenland Ice Sheet: A response to global warming, *J. Clim.*, *21*(2), 331–341.
- Hanna, E., X. Fettweis, S. H. Mernild, J. Cappelien, M. H. Ribergaard, C. A. Shuman, K. Steffen, L. Wood, and T. L. Mote (2014), Atmospheric and oceanic climate forcing of the exceptional Greenland ice sheet surface melt in summer 2012, *Int. J. Climatol.*, *34*(4), 1022–1037.
- Harding, R. J., A. J. Dolman, D. Gerten, I. Haddeland, C. Prudhomme, and P. van Oevelen (2014), Data, models and uncertainties in the global water cycle, in *The Global Water System in the Anthropocene*, edited by A. Bhaduri et al., pp. 45–58, Springer, Cham, Switzerland.
- Hasholt, B., N. Bobrovitskaya, J. Bogen, J. McNamara, S. H. Mernild, D. Milburn, and D. Walling (2006), Sediment transport to the Arctic Ocean and adjoining cold oceans, *Nord. Hydrol.*, *37*(4–5), 413–432.
- Helm, V., A. Humbert, and H. Miller (2014), Elevation and elevation change of Greenland and Antarctica derived from CryoSat-2, *Cryosphere*, *8*(4), 1539–1559.
- Hinzman, L. D., et al. (2005), Evidence and Implications of recent climate change in Northern Alaska and other Arctic Regions, *Clim. Change*, *72*(3), 251–298, doi:10.1007/s10584-005-5352-2.
- Hinzman, L. D., C. J. Deal, A. D. McGuire, S. H. Mernild, I. V. Polyakov, and J. E. Walsh (2013), Trajectory of the Arctic as an integrated system, *Ecol. Appl.*, *23*(8), 1837–1868, doi:10.1890/11-1498.1.
- Holland, M. M., J. Finnis, A. P. Barrett, and M. C. Serreze (2007), Projected changes in Arctic Ocean freshwater budgets, *J. Geophys. Res.*, *112*, G04555, doi:10.1029/2006JG000354.
- Holmes, R. M., B. J. Peterson, V. V. Gordeev, A. V. Zhulidov, M. Meybeck, R. B. Lammers, and C. J. Vörösmarty (2000), Flux of nutrients from Russian rivers to the Arctic Ocean: Can we establish a baseline against which to judge future changes?, *Water Resour. Res.*, *36*, 2309–2320, doi:10.1029/2000WR900099.

- Holmes, R. M., et al. (2011), Seasonal and annual fluxes of nutrients and organic matter from large rivers to the Arctic Ocean and surrounding seas, *Estuaries Coasts*, 35(2), 369–382, doi:10.1007/s12237-011-9386-6.
- Holmes, R. M., et al. (2012), Seasonal and annual fluxes of nutrients and organic matter from large rivers to the Arctic Ocean and surrounding seas, *Estuaries Coasts*, 35(2), 369–382, doi:10.1007/s12237-011-9386-6.
- Holmes, R. M., M. T. Coe, G. J. Fiske, T. Gurtovaya, J. W. McClelland, A. I. Shiklomanov, R. G. Spencer, S. E. Tank, and A. V. Zhulidov (2013), Climate change impacts on the hydrology and biogeochemistry of Arctic rivers, in *Climatic Change and Global Warming of Inland Waters: Impacts and Mitigation for Ecosystems and Societies*, edited by C. R. Goldman, M. Kumagai, and R. D. Robarts, pp. 3–26, John Wiley, Hoboken, N. J.
- Hu, F. S., P. E. Higuera, P. Duffy, M. L. Chipman, A. V. Rocha, A. M. Young, R. Kelly, and M. C. Dietze (2015), Arctic tundra fires: Natural variability and responses to climate change, *Front. Ecol. Environ.*, 13(7), 369–377.
- Iijima, Y., A. N. Fedorov, H. Park, K. Suzuki, H. Yabuki, T. C. Maximov, and T. Ohata (2010), Abrupt increases in soil temperatures following increased precipitation in a permafrost region, central Lena River basin, Russia, *Permafrost Periglacial Processes*, 21(1), 30–41.
- Iijima, Y., T. Ohta, A. Kotani, A. N. Fedorov, Y. Kodama, and T. C. Maximov (2014), Sap flow changes in relation to permafrost degradation under increasing precipitation in an eastern Siberian larch forest, *Ecohydrology*, 7(2), 177–187.
- Instanes, A., V. Kokorev, R. Janowicz, O. Bruland, K. Sand, and T. D. Prowse (2016), Changes to freshwater systems affecting Arctic infrastructure and natural resources, *J. Geophys. Res. Biogeosci.*, 121, doi:10.1002/2015JG003125.
- Jaramillo, F., and G. Destouni (2014), Developing water change spectra and distinguishing change drivers worldwide, *Geophys. Res. Lett.*, 41, 8377–8386, doi:10.1002/2014GL061848.
- Jaramillo, F., and G. Destouni (2015), Local flow regulation and irrigation raise global human water consumption and footprint, *Science*, 350(6265), 1248–1251, doi:10.1126/science.aad1010.
- Jarsjö, J., S. M. Asokan, C. Prieto, A. Bring, and G. Destouni (2012), Hydrological responses to climate change conditioned by historic alterations of land-use and water-use, *Hydrol. Earth Syst. Sci.*, 16(5), 1335–1347, doi:10.5194/hess-16-1335-2012.
- Jepsen, S. M., C. I. Voss, M. A. Walvoord, B. J. Minsley, and J. Rover (2013), Linkages between lake shrinkage/expansion and sublacustrine permafrost distribution determined from remote sensing of interior Alaska, USA, *Geophys. Res. Lett.*, 40, 882–887, doi:10.1002/grl.50187.
- Jones, B. M., G. Grosse, C. D. Arp, M. C. Jones, K. M. Walter Anthony, and V. E. Romanovsky (2011), Modern thermokarst lake dynamics in the continuous permafrost zone, northern Seward Peninsula, Alaska, *J. Geophys. Res.*, 116, G00M03, doi:10.1029/2011JG001666.
- Jorgenson, M. T., Y. L. Shur, and T. E. Osterkamp (2008), Thermokarst in Alaska, in *Proceedings of the Ninth International Conference on Permafrost*, vol. 29, pp. 869–876, Inst. of North. Eng., Univ. of Alaska, Fairbanks, Alaska.
- Kane, D. L. (1981), Physical mechanics of aufeis growth, *Can. J. Civ. Eng.*, 8(2), 186–195.
- Kane, D. L. (1997), The impact of hydrologic perturbations on arctic ecosystems induced by climate change, in *Global Change and Arctic Terrestrial Ecosystems*, edited by W. C. Oechel et al., pp. 63–81, Springer, New York.
- Kane, D. L., and S. L. Stuefer (2015), Reflecting on the status of precipitation data collection in Alaska: A case study, *Hydrol. Res.*, 46, 478–493.
- Kane, D. L., L. Hinzman, R. Gieck, J. McNamara, E. Youcha, and J. Oatley (2008), Contrasting extreme runoff events in areas of continuous permafrost, Arctic Alaska, *Hydrol. Res.*, 39, doi:10.2166/nh.2008.005.
- Karlsson, J. M., A. Bring, G. D. Peterson, L. J. Gordon, and G. Destouni (2011), Opportunities and limitations to detect climate-related regime shifts in inland Arctic ecosystems through eco-hydrological monitoring, *Environ. Res. Lett.*, 6(1), 014015, doi:10.1088/1748-9326/6/1/014015.
- Karlsson, J. M., S. W. Lyon, and G. Destouni (2012), Thermokarst lake, hydrological flow and water balance indicators of permafrost change in western Siberia, *J. Hydrol.*, 464–465, 459–466, doi:10.1016/j.jhydrol.2012.07.037.
- Karlsson, J. M., F. Jaramillo, and G. Destouni (2015), Hydro-climatic and lake change patterns in Arctic permafrost and non-permafrost areas, *J. Hydrol.*, 529(Part 1), 134–145, doi:10.1016/j.jhydrol.2015.07.005.
- Kaser, G., J. G. Cogley, M. B. Dyurgerov, M. F. Meier, and A. Ohmura (2006), Mass balance of glaciers and ice caps: Consensus estimates for 1961–2004, *Geophys. Res. Lett.*, 33, L19501, doi:10.1029/2006GL027511.
- Kattsov, V. M., et al. (2005), Future climate change: Modeling and scenarios for the Arctic, in *Arctic Climate Impact Assessment*, pp. 99–150, Cambridge Univ. Press, Cambridge, U. K.
- Kattsov, V. M., J. E. Walsh, W. L. Chapman, V. A. Govorkova, T. V. Pavlova, and X. Zhang (2007), Simulation and projection of Arctic freshwater budget components by the IPCC AR4 global climate models, *J. Hydrometeorol.*, 8(3), 571–589.
- Kharin, V. V., F. W. Zwiers, X. Zhang, and M. Wehner (2013), Changes in temperature and precipitation extremes in the CMIP5 ensemble, *Clim. Change*, 119(2), 345–357.
- Kirkham, R. V., L. B. Chorlton, and J. J. Carriere (1995), Generalized geology of the world, in *Generalized Geological Map of the World and Linked Databases*, Open File 2915d, Geol. Surv. of Canada.
- Koch, J. C., R. L. Runkel, R. Striegl, and D. M. McKnight (2013), Hydrologic controls on the transport and cycling of carbon and nitrogen in a boreal catchment underlain by continuous permafrost, *J. Geophys. Res. Biogeosci.*, 118, 698–712, doi:10.1002/jgrg.20058.
- Koirala, S., Y. Hirabayashi, R. Mahendran, and S. Kanae (2014), Global assessment of agreement among streamflow projections using CMIP5 model outputs, *Environ. Res. Lett.*, 9(6), 064017, doi:10.1088/1748-9326/9/6/064017.
- Koven, C. D., W. J. Riley, and A. Stern (2013), Analysis of permafrost thermal dynamics and response to climate change in the CMIP5 Earth System Models, *J. Clim.*, 26, doi:10.1175/JCLI-D-12-00228.1.
- Krasting, J. P., A. J. Broccoli, K. W. Dixon, and J. R. Lanzante (2013), Future changes in Northern Hemisphere snowfall, *J. Clim.*, 26(20), 7813–7828.
- Lainé, A., H. Nakamura, K. Nishii, and T. Miyasaka (2014), A diagnostic study of future evaporation changes projected in CMIP5 climate models, *Clim. Dyn.*, 42(9–10), 2745–2761.
- Lammers, R. B., A. I. Shiklomanov, C. J. Vörösmarty, B. M. Fekete, and B. J. Peterson (2001), Assessment of contemporary Arctic river runoff based on observational discharge records, *J. Geophys. Res.*, 106, 3321–3334, doi:10.1029/2000JD900444.
- Langer, M., S. Westermann, S. Muster, K. Piel, and J. Boike (2011), The surface energy balance of a polygonal tundra site in northern Siberia. Part 2: Winter, *Cryosphere*, 5, 509–524.
- Latifovic, R., and D. Pouliot (2007), Analysis of climate change impacts on lake ice phenology in Canada using the historical satellite data record, *Remote Sens. Environ.*, 106, 492–507.
- Leconte, R., M. Temimi, N. Chaouch, F. Brissette, and T. Toussaint (2008), On the use of satellite passive microwave data for estimating surface soil wetness in the Mackenzie River Basin, in *Cold Region Atmospheric and Hydrologic Studies. The Mackenzie GEWEX Experience*, edited by M.-K. Woo, pp. 59–79, Springer, Berlin.
- Lehner, B., and P. Döll (2004), Development and validation of a global database of lakes, reservoirs and wetlands, *J. Hydrol.*, 296(1), 1–22.
- Lettenmaier, D. P., D. Alsdorf, J. Dozier, G. J. Huffman, M. Pan, and E. F. Wood (2015), Inroads of remote sensing into hydrologic science during the WRR era, *Water Resour. Res.*, 51, 7309–7342, doi:10.1002/2015WR017616.

- Lique, C., M. M. Holland, Y. Dibike, D. M. Lawrence, and J. Screen (2016), Modeling the Arctic freshwater system and its integration in the global system: Lessons learned and future challenges, *J. Geophys. Res. Biogeosci.*, *121*, doi:10.1002/2015JG003120.
- Liston, G. E., and C. A. Hiemstra (2011), The changing cryosphere: Pan-Arctic snow trends (1979–2009), *J. Clim.*, *24*(21), 5691–5712, doi:10.1175/JCLI-D-11-00081.1.
- Liston, G. E., and S. H. Mernild (2012), Greenland freshwater runoff. Part I: A runoff routing model for glaciated and nonglaciated landscapes (HydroFlow), *J. Clim.*, *25*(17), 5997–6014, doi:10.1175/JCLI-D-11-00591.1.
- Livingstone, D. M., R. Adrian, T. Blenckner, G. George, and G. A. Weyhenmeyer (2010), Lake ice phenology, in *The Impact of Climate Change on European Lakes*, edited by G. D. George, pp. 51–61, Springer, Dordrecht, Netherlands.
- Lyon, S. W., and G. Destouni (2010), Changes in catchment-scale recession flow properties in response to permafrost thawing in the Yukon River Basin, *Int. J. Climatol.*, *30*(14), 2138–2145, doi:10.1002/joc.1993.
- Lyon, S. W., G. Destouni, R. Giesler, C. Humborg, M. Mörth, J. Seibert, J. Karlsson, and P. A. Troch (2009), Estimation of permafrost thawing rates in a sub-arctic catchment using recession flow analysis, *Hydrol. Earth Syst. Sci.*, *13*(5), 595–604.
- Magnuson, J. J., et al. (2000), Historical trends in lake and river ice cover in the Northern Hemisphere, *Science*, *289*(5485), 1743–1746.
- Mahlstein, I., R. W. Portmann, J. S. Daniel, S. Solomon, and R. Knutti (2012), Perceptible changes in regional precipitation in a future climate, *Geophys. Res. Lett.*, *39*, L05701, doi:10.1029/2011GL050738.
- Mantua, N. J., and S. R. Hare (2002), The Pacific decadal oscillation, *J. Oceanogr.*, *58*(1), 35–44.
- Marsh, P., and S. C. Bigras (1988), Evaporation from Mackenzie delta lakes, NWT, Canada, *Arct. Alp. Res.*, *20*, 220–229.
- Mazi, K., A. D. Koussis, and G. Destouni (2013), Tipping points for seawater intrusion in coastal aquifers under rising sea level, *Environ. Res. Lett.*, *8*(1), 014001.
- McClelland, J. W., R. M. Holmes, B. J. Peterson, and M. Stieglitz (2004), Increasing river discharge in the Eurasian Arctic: Consideration of dams, permafrost thaw, and fires as potential agents of change, *J. Geophys. Res.*, *109*, D18102, doi:10.1029/2004JD004583.
- McClelland, J. W., S. J. Déry, B. J. Peterson, R. M. Holmes, and E. F. Wood (2006), A pan-arctic evaluation of changes in river discharge during the latter half of the 20th century, *Geophys. Res. Lett.*, *33*, L06715, doi:10.1029/2006GL025753.
- McClelland, J. W., et al. (2008), Development of a Pan-Arctic database for river chemistry, *Eos Trans. AGU*, *89*(24), 217–218, doi:10.1029/2008EO240001.
- McClelland, J. W., S. E. Tank, R. G. M. Spencer, and A. I. Shiklomanov (2015), Coordination and sustainability of river observing activities in the Arctic, *Arctic*, *68*(5), 1–10.
- Mernild, S. H., and G. E. Liston (2012), Greenland freshwater runoff. Part II: Distribution and trends, 1960–2010, *J. Clim.*, *25*(17), 6015–6035.
- Mernild, S. H., J. K. Malmros, J. C. Yde, and N. T. Knudsen (2012), Multi-decadal marine- and land-terminating glacier recession in the Ammassalik region, southeast Greenland, *Cryosphere*, *6*(3), 625–639.
- Mernild, S. H., G. E. Liston, and C. A. Hiemstra (2014), Northern Hemisphere glacier and ice cap surface mass balance and contribution to sea level rise, *J. Clim.*, *27*(15), 6051–6073.
- Mernild, S. H., E. Hanna, J. R. McConnell, M. Sigl, A. P. Beckerman, J. C. Yde, J. Cappelen, J. K. Malmros, and K. Steffen (2015), Greenland precipitation trends in a long-term instrumental climate context (1890–2012): Evaluation of coastal and ice core records, *Int. J. Climatol.*, *35*, 303–320, doi:10.1002/joc.3986.
- Morgenstern, A., G. Grosse, F. Günther, I. Fedorova, and L. Schirmer (2011), Spatial analyses of thermokarst lakes and basins in Yedoma landscapes of the Lena Delta, *Cryosphere Discuss.*, *5*, 1495–1545.
- Mountain Research Initiative EDW Working Group (2015), Elevation-dependent warming in mountain regions of the world, *Nat. Clim. Change*, *5*(5), 424–430, doi:10.1038/nclimate2563.
- Mu, Q., L. A. Jones, J. S. Kimball, K. C. McDonald, and S. W. Running (2009), Satellite assessment of land surface evapotranspiration for the pan-Arctic domain, *Water Resour. Res.*, *45*, W09420, doi:10.1029/2008WR007189.
- Mueller, B., et al. (2011), Evaluation of global observations-based evapotranspiration datasets and IPCC AR4 simulations, *Geophys. Res. Lett.*, *38*, L06402, doi:10.1029/2010GL046230.
- Nghiem, S. V., D. K. Hall, T. L. Mote, M. Tedesco, M. R. Albert, K. Keegan, C. A. Shuman, N. E. DiGirolamo, and G. Neumann (2012), The extreme melt across the Greenland ice sheet in 2012, *Geophys. Res. Lett.*, *39*, L20502, doi:10.1029/2012GL053611.
- Nilsson, L. M., G. Destouni, J. Berner, A. A. Dudarev, G. Mulvad, J. Ø. Odland, A. Parkinson, C. Tikhonov, A. Rautio, and B. Evengård (2013), A call for urgent monitoring of food and water security based on relevant indicators for the Arctic, *Ambio*, *42*(7), 816–822.
- Nowak, A., and A. Hodson (2013), Hydrological response of a High-Arctic catchment to changing climate over the past 35 years: A case study of Bayelva watershed, Svalbard, *Polar Res.*, *32*, 19,691, doi:10.3402/polar.v32i0.19691.
- Olson, D. M., et al. (2001), Terrestrial ecoregions of the world: A new map of life on Earth, *BioScience*, *51*(11), 933–938.
- Overduin, P. P., M. C. Strzelecki, M. N. Grigoriev, N. Couture, H. Lantuit, D. St-Hilaire-Gravel, F. Günther, and S. Wetterich (2014), Coastal changes in the Arctic, *Geol. Soc. London Spec. Publ.*, *388*(1), 103–129.
- Overeem, I., and J. P. M. Syvitski (2010), Shifting discharge peaks in Arctic rivers, 1977–2007, *Geogr. Ann. Ser. Phys. Geogr.*, *92*(2), 285–296.
- Parent, M. B., and D. Verbyla (2010), The browning of Alaska's boreal forest, *Remote Sens.*, *2*(12), 2729–2747.
- Park, H., J. Walsh, A. N. Fedorov, A. B. Sherstiukov, Y. Iijima, and T. Ohata (2013), The influence of climate and hydrological variables on opposite anomaly in active-layer thickness between Eurasian and North American watersheds, *Cryosphere*, *7*(2), 631–645.
- Paul, F., and F. Svoboda (2010), A new glacier inventory on southern Baffin Island, Canada, from ASTER data: II. Data analysis, glacier change and applications, *Ann. Glaciol.*, *50*(53), 22–31.
- Pavelsky, T. M., and L. C. Smith (2006), Intercomparison of four global precipitation data sets and their correlation with increased Eurasian river discharge to the Arctic Ocean, *J. Geophys. Res.*, *111*, D21112, doi:10.1029/2006JD007230.
- Pearson, R. G., S. J. Phillips, M. M. Lorant, P. S. A. Beck, T. Damoulas, S. J. Knight, and S. J. Goetz (2013), Shifts in Arctic vegetation and associated feedbacks under climate change, *Nat. Clim. Change*, *3*(7), 673–677, doi:10.1038/nclimate1858.
- Peterson, B. J., R. M. Holmes, J. W. McClelland, C. J. Vörösmarty, R. B. Lammers, A. I. Shiklomanov, I. A. Shiklomanov, and S. Rahmstorf (2002), Increasing river discharge to the Arctic Ocean, *Science*, *298*, 2171–2173.
- Peterson, B. J., J. McClelland, R. Curry, R. M. Holmes, J. E. Walsh, and K. Aagaard (2006), Trajectory shifts in the Arctic and subarctic freshwater cycle, *Science*, *313*(5790), 1061–1066, doi:10.1126/science.1122593.
- Prowse, T., R. Shrestha, B. Bonsal, and Y. Dibike (2010), Changing spring air-temperature gradients along large northern rivers: Implications for severity of river-ice floods, *Geophys. Res. Lett.*, *37*, L19706, doi:10.1029/2010GL044878.
- Prowse, T., K. Alfredsen, S. Beltaos, B. Bonsal, C. Duguay, A. Korhola, J. McNamara, W. F. Vincent, V. Vuglinsky, and G. A. Weyhenmeyer (2011), Arctic freshwater ice and its climatic role, *Ambio*, *40*(1), 46–52.
- Prowse, T., A. Bring, J. Mård, and E. Carmack (2015a), Arctic Freshwater Synthesis: Introduction, *J. Geophys. Res. Biogeosci.*, *120*, 2121–2131, doi:10.1002/2015JG003127.

- Prowse, T., A. Bring, J. M. Karlsson, E. Carmack, M. Holland, A. Instanes, T. Vihma, and F. J. Wrona (2015b), Arctic Freshwater Synthesis: Summary of key emerging issues, *J. Geophys. Res. Biogeosci.*, *120*, 1887–1893, doi:10.1002/2015JG003128.
- Prowse, T. D., and K. Brown (2010), Hydro-ecological effects of changing Arctic river and lake ice covers: A review, *Hydrol. Res.*, *41*(6), 454–461.
- Prowse, T. D., et al. (2005), Terrestrial Cryospheric and Hydrologic Processes and Systems, International Conference on Arctic Research Planning, Sci. Plan. Work. Group 7, Copenhagen, Denmark.
- Quin, A., F. Jaramillo, and G. Destouni (2015), Dissecting the ecosystem service of large-scale pollutant retention: The role of wetlands and other landscape features, *Ambio*, *44*(1), 127–137, doi:10.1007/s13280-014-0594-8.
- Quinton, W. L., T. Shirazi, S. K. Carey, and J. W. Pomeroy (2005), Soil water storage and active-layer development in a sub-alpine tundra hillslope, southern Yukon Territory, Canada, *Permafrost Periglacial Processes*, *16*(4), 369–382.
- Quinton, W. L., M. Hayashi, and L. E. Chasmer (2011), Permafrost-thaw-induced land-cover change in the Canadian subarctic: Implications for water resources, *Hydrol. Processes*, *25*(1), 152–158.
- Rachold, V., A. Alabyan, H.-W. Hubberten, V. N. Korotaev, and A. A. Zaitsev (1996), Sediment transport to the Laptev Sea—Hydrology and geochemistry of the Lena River, *Polar Res.*, *15*(2), 183–196.
- Rachold, V., M. N. Grigoriev, F. E. Are, S. Solomon, E. Reimnitz, H. Kassens, and M. Antonow (2000), Coastal erosion vs riverine sediment discharge in the Arctic Shelf seas, *Int. J. Earth Sci.*, *89*(3), 450–460.
- Rahmstorf, S., et al. (2005), Thermohaline circulation hysteresis: A model intercomparison, *Geophys. Res. Lett.*, *32*, L23605, doi:10.1029/2005GL023655.
- Rahmstorf, S., J. E. Box, G. Feulner, M. E. Mann, A. Robinson, S. Rutherford, and E. J. Schaffernicht (2015), Exceptional twentieth-century slowdown in Atlantic Ocean overturning circulation, *Nat. Clim. Change*, *5*, 475–480.
- Räisänen, J. (2008), Warmer climate: Less or more snow?, *Clim. Dyn.*, *30*(2–3), 307–319.
- Rawlins, M. A., M. C. Serreze, R. Schroeder, X. Zhang, and K. C. McDonald (2009), Diagnosis of the record discharge of Arctic-draining Eurasian rivers in 2007, *Environ. Res. Lett.*, *4*(4), 045011, doi:10.1088/1748-9326/4/4/045011.
- Rawlins, M. A., et al. (2010), Analysis of the Arctic system for freshwater cycle intensification: Observations and expectations, *J. Clim.*, *23*(21), 5715–5737, doi:10.1175/2010JCLI3421.1.
- Raymond, P. A., J. W. McClelland, R. M. Holmes, A. V. Zhulidov, K. Mull, B. J. Peterson, R. G. Striegl, G. R. Aiken, and T. Y. Gurtovaya (2007), Flux and age of dissolved organic carbon exported to the Arctic Ocean: A carbon isotopic study of the five largest arctic rivers, *Global Biogeochem. Cycles*, *21*, GB4011, doi:10.1029/2007GB002934.
- Raymond, P. A., et al. (2013), Global carbon dioxide emissions from inland waters, *Nature*, *503*(7476), 355–359, doi:10.1038/nature12760.
- Rennermalm, A. K., E. F. Wood, and T. J. Troy (2010), Observed changes in pan-arctic cold-season minimum monthly river discharge, *Clim. Dyn.*, *35*(6), 923–939.
- Rennermalm, A. K., A. Bring, and T. L. Mote (2012), Spatial and scale-dependent controls on North American Pan-Arctic minimum river discharge, *Geogr. Anal.*, *44*(3), 202–218, doi:10.1111/j.1538-4632.2012.00849.x.
- Rouse, W. R., C. J. Oswald, J. Binyamin, C. Spence, W. M. Schertzer, P. D. Blanken, N. Bussi eres, and C. R. Duguay (2005), The role of northern lakes in a regional energy balance, *J. Hydrometeorol.*, *6*(3), 291–305.
- Rouse, W. R., P. D. Blanken, C. R. Duguay, C. J. Oswald, and W. M. Schertzer (2008), Climate-lake interactions, in *Cold Region Atmospheric and Hydrologic Studies. The Mackenzie GEWEX Experience*, pp. 139–160, Springer, Heidelberg, Germany.
- Rovensek, R. J., L. D. Hinzman, and D. L. Kane (1996), Hydrology of a tundra wetland complex on the Alaskan Arctic Coastal Plain, USA, *Arct. Alp. Res.*, *28*, 311–317.
- Rowland, J. C., et al. (2010), Arctic landscapes in transition: Responses to thawing permafrost, *Eos Trans. AGU*, *91*(26), 229–230, doi:10.1029/2010EO260001.
- Ruhoff, A. L., A. R. Paz, L. Aragao, Q. Mu, Y. Malhi, W. Collischonn, H. R. Rocha, and S. W. Running (2013), Assessment of the MODIS global evapotranspiration algorithm using eddy covariance measurements and hydrological modelling in the Rio Grande basin, *Hydrol. Sci. J.*, *58*(8), 1658–1676.
- Sauchyn, D. J., E. M. Barrow, R. Hopkinson, and P. R. Leavitt (2002), Aridity on the Canadian plains, *G ogr. Phys. Quat.*, *56*(2–3), 247, doi:10.7202/009109ar.
- Schindler, D. W., and W. F. Donahue (2006), An impending water crisis in Canada's western prairie provinces, *Proc. Natl. Acad. Sci. U.S.A.*, *103*(19), 7210–7216, doi:10.1073/pnas.0601568103.
- Schuur, E. A. G., et al. (2015), Climate change and the permafrost carbon feedback, *Nature*, *520*(7546), 171–179.
- Sellers, P., M. Rienecker, S. Frolking, and D. Randall (2012), *Earth System Modeling and Field Experiments in the Arctic–Boreal Zone, NASA Workshop Rep.*, NASA, Hanover, Md.
- Semenova, O., L. Lebedeva, and Y. Vinogradov (2013), Simulation of subsurface heat and water dynamics, and runoff generation in mountainous permafrost conditions, in the Upper Kolyma River basin, Russia, *Hydrogeol. J.*, *21*(1), 107–119.
- Semenova, O., L. Lebedeva, N. Volkova, I. Korenev, M. Forkel, J. Eberle, and M. Urban (2015), Detecting immediate wildfire impact on runoff in a poorly-gauged mountainous permafrost basin, *Hydrol. Sci. J.*, *60*(7–8), 1225–1241, doi:10.1080/02626667.2014.959960.
- Serreze, M. C., and C. M. Hurst (2000), Representation of mean Arctic precipitation from NCEP-NCAR and ERA reanalyses, *J. Clim.*, *13*(1), 182–201.
- Serreze, M. C., A. P. Barrett, A. G. Slater, R. A. Woodgate, K. Aagaard, R. B. Lammers, M. Steele, R. Moritz, M. Meredith, and C. M. Lee (2006), The large-scale freshwater cycle of the Arctic, *J. Geophys. Res.*, *111*, C11010, doi:10.1029/2005JC003424.
- Shahgedanova, M., G. Nosenko, I. Bushueva, and M. Ivanov (2012), Changes in area and geodetic mass balance of small glaciers, Polar Urals, Russia, 1950–2008, *J. Glaciol.*, *58*(211), 953–964.
- Shaw, D. A., G. Vanderkamp, F. M. Conly, A. Pietroniro, and L. Martz (2012), The fill–spill hydrology of prairie wetland complexes during drought and deluge, *Hydrol. Processes*, *26*(20), 3147–3156, doi:10.1002/hyp.8390.
- Sheng, Y., L. C. Smith, G. M. MacDonald, K. V. Kremenetski, K. E. Frey, A. A. Velichko, M. Lee, D. W. Beilman, and P. Dubinin (2004), A high-resolution GIS-based inventory of the west Siberian peat carbon pool, *Global Biogeochem. Cycles*, *18*, GB3004, doi:10.1029/2003GB002190.
- Shiklomanov, A. I., and R. B. Lammers (2009), Record Russian river discharge in 2007 and the limits of analysis, *Environ. Res. Lett.*, *4*(4), 045015, doi:10.1088/1748-9326/4/4/045015.
- Shiklomanov, A. I., R. B. Lammers, and C. J. V or smarty (2002), Widespread decline in hydrological monitoring threatens pan-Arctic research, *Eos Trans. AGU*, *83*, 13–17, doi:10.1029/2002EO000007.
- Shiklomanov, A. I., T. I. Yakovleva, R. B. Lammers, I. P. Karasev, C. J. V or smarty, and E. Linder (2006), Cold region river discharge uncertainty—Estimates from large Russian rivers, *J. Hydrol.*, *326*, 231–256.
- Shiklomanov, A. I., R. B. Lammers, M. A. Rawlins, L. C. Smith, and T. M. Pavelsky (2007), Temporal and spatial variations in maximum river discharge from a new Russian data set, *J. Geophys. Res.*, *112*, G04553, doi:10.1029/2006JG000352.

- Sillmann, J., V. V. Kharin, F. W. Zwiers, X. Zhang, and D. Bronaugh (2013), Climate extremes indices in the CMIP5 multimodel ensemble: Part 2. Future climate projections, *J. Geophys. Res. Atmos.*, *118*, 2473–2493, doi:10.1002/jgrd.50188.
- Sjöberg, Y., A. Frampton, and S. W. Lyon (2013), Using streamflow characteristics to explore permafrost thawing in northern Swedish catchments, *Hydrogeol. J.*, *21*(1), 121–131, doi:10.1007/s10040-012-0932-5.
- Slater, A. G., and D. M. Lawrence (2013), Diagnosing present and future permafrost from climate models, *J. Clim.*, *26*(15), 5608–5623.
- Slaughter, C. W., J. W. Hilgert, and E. H. Culp (1983), Summer streamflow and sediment yield from discontinuous-permafrost headwaters catchments, in *Proceedings, Fourth International Conference on Permafrost*, pp. 1172–1177, Natl. Acad. Press, Washington, D. C.
- Smith, K. B., C. E. Smith, S. F. Forest, and A. J. Richard (2007), *A Field Guide to the Wetlands of the Boreal Plains Ecozone of Canada*, Ducks Unlimited Canada, West Boreal Off., Edmonton, Alberta.
- Smith, L. C., and T. M. Pavelsky (2008), Estimation of river discharge, propagation speed, and hydraulic geometry from space: Lena River, Siberia, *Water Resour. Res.*, *44*, W03427, doi:10.1029/2007WR006133.
- Smith, L. C., Y. Sheng, G. M. MacDonald, and L. D. Hinzman (2005), Disappearing Arctic lakes, *Science*, *308*(5727), 1429, doi:10.1126/science.1108142.
- Smith, L. C., T. M. Pavelsky, G. M. MacDonald, A. I. Shiklomanov, and R. B. Lammers (2007), Rising minimum daily flows in northern Eurasian rivers: A growing influence of groundwater in the high-latitude hydrologic cycle, *J. Geophys. Res.*, *112*, G04S47, doi:10.1029/2006JG000327.
- Smith, L. C., et al. (2015), Efficient meltwater drainage through supraglacial streams and rivers on the southwest Greenland ice sheet, *Proc. Natl. Acad. Sci. U.S.A.*, *112*(4), 1001–1006, doi:10.1073/pnas.1413024112.
- Spence, C., and M. Woo (2002), Hydrology of subarctic Canadian Shield: Bedrock upland, *J. Hydrol.*, *262*(1), 111–127.
- Spence, C., and M. Woo (2003), Hydrology of subarctic Canadian shield: Soil-filled valleys, *J. Hydrol.*, *279*(1), 151–166.
- Spence, C., S. V. Kokelj, S. A. Kokelj, M. McKluskie, and N. Hedstrom (2014), Evidence of a change in water chemistry in Canada's subarctic associated with enhanced winter streamflow, *J. Geophys. Res. Biogeosci.*, *120*, 113–127, doi:10.1002/2014JG002809.
- St. Jacques, J.-M., and D. J. Sauchyn (2009), Increasing winter baseflow and mean annual streamflow from possible permafrost thawing in the Northwest Territories, Canada, *Geophys. Res. Lett.*, *36*, L01401, doi:10.1029/2008GL035822.
- Stocks, B. J., et al. (1998), Climate change and forest fire potential in Russian and Canadian boreal forests, *Clim. Change*, *38*(1), 1–13.
- Strandmark, A., A. Bring, S. A. O. Cousins, G. Destouni, H. Kautsky, G. Kolb, M. de la Torre Castro, and P. A. Hambäck (2015), Climate change effects in the borderland between land and sea: An overseen issue, *Ambio*, *44*(Suppl. 1), 28–38, doi:10.1007/s13280-014-0586-8.
- Stuefer, S., D. Yang, and A. Shiklomanov (2011), Effect of streamflow regulation on mean annual discharge variability of the Yenisei River, in *Cold Regions Hydrology in a Changing Climate*, edited by D. Yang, P. Marsh, and A. Gelfan, pp. 27–32, International Association of Hydrological Sciences, Wallingford, U. K.
- Stuefer, S., D. L. Kane, and G. E. Liston (2013), In situ snow water equivalent observations in the US Arctic, *Hydrol. Res.*, *44*(1), 21–34.
- Sturm, M., J. Holmgren, J. P. McFadden, G. E. Liston, F. S. Chapin, and C. H. Racine (2001), Snow-shrub interactions in arctic tundra: A hypothesis with climatic implications, *J. Clim.*, *14*(3), 336–344.
- Sturm, M., T. Douglas, C. Racine, and G. E. Liston (2005), Changing snow and shrub conditions affect albedo with global implications, *J. Geophys. Res.*, *110*, G01004, doi:10.1029/2005JG000013.
- Su, F., J. C. Adam, L. C. Bowling, and D. P. Lettenmaier (2005), Streamflow simulations of the terrestrial Arctic domain, *J. Geophys. Res.*, *110*, D08112, doi:10.1029/2004JD005518.
- Sun, G., A. Noormets, J. Chen, and S. G. McNulty (2008), Evapotranspiration estimates from eddy covariance towers and hydrologic modeling in managed forests in Northern Wisconsin, USA, *Agric. For. Meteorol.*, *148*(2), 257–267.
- Syvitski, J. P., et al. (2005), Dynamics of the coastal zone, in *Coastal Fluxes in the Anthropocene*, pp. 39–94, Springer, Heidelberg, Germany.
- Tan, A., J. C. Adam, and D. P. Lettenmaier (2011), Change in spring snowmelt timing in Eurasian Arctic rivers, *J. Geophys. Res.*, *116*, D03101, doi:10.1029/2010JD014337.
- Tank, S. E., P. A. Raymond, R. G. Striegl, J. W. McClelland, R. M. Holmes, G. J. Fiske, and B. J. Peterson (2012), A land-to-ocean perspective on the magnitude, source and implication of DIC flux from major Arctic rivers to the Arctic Ocean, *Global Biogeochem. Cycles*, *26*, GB4018, doi:10.1029/2011GB004192.
- Tape, K., M. Sturm, and C. Racine (2006), The evidence for shrub expansion in Northern Alaska and the Pan-Arctic, *Global Change Biol.*, *12*(4), 686–702.
- Tape, K., J. M. Hallinger, J. M. Welker, and R. W. Ruess (2012), Landscape heterogeneity of shrub expansion in Arctic Alaska, *Ecosystems*, *15*, 711–724.
- Tetzlaff, D., J. Buttle, S. K. Carey, K. McGuire, H. Laudon, and C. Soulsby (2015), Tracer-based assessment of flow paths, storage and runoff generation in northern catchments: A review, *Hydrol. Processes*, *29*, 3475–3490, doi:10.1002/hyp.10412.
- Törnqvist, R., J. Jarsjö, J. Pietroń, A. Bring, P. Rogberg, S. M. Asokan, and G. Destouni (2014), Evolution of the hydro-climate system in the Lake Baikal basin, *J. Hydrol.*, *519*, 1953–1962, doi:10.1016/j.jhydrol.2014.09.074.
- Troy, T. J., J. Sheffield, and E. F. Wood (2012), The role of winter precipitation and temperature on northern Eurasian streamflow trends, *J. Geophys. Res.*, *117*, D05131, doi:10.1029/2011JD016208.
- Turcotte, B., B. Morse, N. E. Bergeron, and A. G. Roy (2011), Sediment transport in ice-affected rivers, *J. Hydrol.*, *409*(1), 561–577.
- Van der Ent, R. J., H. H. Savenije, B. Schaefli, and S. C. Steele-Dunne (2010), Origin and fate of atmospheric moisture over continents, *Water Resour. Res.*, *46*, W09525, doi:10.1029/2010WR009127.
- Van der Velde, Y., N. Vercauteren, F. Jaramillo, S. C. Dekker, G. Destouni, and S. W. Lyon (2014), Exploring hydroclimatic change disparity via the Budyko framework, *Hydrol. Processes*, *28*(13), 4110–4118, doi:10.1002/hyp.9949.
- Van Vliet, M. T. H., W. H. P. Franssen, J. R. Yearsley, F. Ludwig, I. Haddeland, D. P. Lettenmaier, and P. Kabat (2013), Global river discharge and water temperature under climate change, *Global Environ. Change*, *23*(2), 450–464, doi:10.1016/j.gloenvcha.2012.11.002.
- Vaughan, D. G., et al. (2013), Observations: Cryosphere, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by T. F. Stocker et al., pp. 317–382, Cambridge Univ. Press, Cambridge, U. K.
- Verbyla, D. (2011), Browning boreal forests of western North America, *Environ. Res. Lett.*, *6*(4), 041003.
- Vernon, C. L., J. L. Bamber, J. E. Box, M. R. Van den Broeke, X. Fettweis, E. Hanna, and P. Huybrechts (2013), Surface mass balance model intercomparison for the Greenland ice sheet, *Cryosphere*, *7*, 599–614.
- Vihma, T., J. Screen, M. Tjernström, B. Newton, X. Zhang, V. Popova, C. Deser, M. Holland, and T. D. Prowse (2016), The atmospheric role in the Arctic water cycle: Processes, past and future changes, and their impacts, *J. Geophys. Res. Biogeosci.*, *121*, doi:10.1002/2015JG003132.
- Walsh, J. E., J. E. Overland, P. Y. Grouisman, and B. Rudolf (2011), Ongoing climate change in the Arctic, *Ambio*, *40*(1), 6–16, doi:10.1007/s13280-011-0211-z.

- Walvoord, M. A., and R. G. Striegl (2007), Increased groundwater to stream discharge from permafrost thawing in the Yukon River basin: Potential impacts on lateral export of carbon and nitrogen, *Geophys. Res. Lett.*, *34*, L12402, doi:10.1029/2007GL030216.
- Walvoord, M. A., C. I. Voss, and T. P. Wellman (2012), Influence of permafrost distribution on groundwater flow in the context of climate-driven permafrost thaw: Example from Yukon Flats Basin, Alaska, United States, *Water Resour. Res.*, *48*, W07524, doi:10.1029/2011WR011595.
- White, D. M., S. Craig Gerlach, P. Loring, A. C. Tidwell, and M. C. Chambers (2007), Food and water security in a changing arctic climate, *Environ. Res. Lett.*, *2*(4), 045018, doi:10.1088/1748-9326/2/4/045018.
- Woo, M. (1986), Permafrost Hydrology in North America¹, *Atmos. Ocean*, *24*(3), 201–234.
- Woo, M., and C. Mielko (2007), An integrated framework of lake-stream connectivity for a semi-arid, subarctic environment, *Hydrol. Processes*, *21*(19), 2668–2674.
- Woo, M.-K., and R. Thorne (2003), Streamflow in the Mackenzie Basin, Canada, *Arctic*, *56*, 328–340.
- Woo, M.-K., and R. Thorne (2014), Winter flows in the Mackenzie drainage system, *Arctic*, *67*(2), 238–256.
- World Glacier Monitoring Service (2013), Glacier Mass Balance Bulletin No. 12 (2010–2011), edited by M. Zemp, 106 pp., ICSU(WDS)/IUGG (IACS)/UNEP/ UNESCO/WMO, World Glacier Monitoring Service, Zurich, Switzerland, doi:10.5904/wgms-fog-2013-11.
- Wrona, F. J., M. Johansson, J. M. Culp, A. Jenkins, J. Mård, I. H. Myers-Smith, T. D. Prowse, W. F. Vincent, and P. A. Wookey (2016), Transitions in Arctic ecosystems: Ecological implications of a changing hydrological regime, *J. Geophys. Res. Biogeosci.*, *121*, doi:10.1002/2015JG003133.
- Yang, D., D. L. Kane, L. D. Hinzman, X. Zhang, T. Zhang, and H. Ye (2002), Siberian Lena River hydrologic regime and recent change, *J. Geophys. Res.*, *107*(D23), 4694, doi:10.1029/2002JD002542.
- Yang, D., B. Ye, and A. Shiklomanov (2004a), Discharge characteristics and changes over the Ob River watershed in Siberia, *J. Hydrometeorol.*, *5*(4), 595–610, doi:10.1175/1525-7541(2004)005<0595:DCACOT>2.0.CO;2.
- Yang, D., B. Ye, and D. L. Kane (2004b), Streamflow changes over Siberian Yenisei river basin, *J. Hydrol.*, *296*(1–4), 59–80, doi:10.1016/j.jhydrol.2004.03.017.
- Yang, D., D. Kane, Z. Zhang, D. Legates, and B. Goodison (2005), Bias corrections of long-term (1973–2004) daily precipitation data over the northern regions, *Geophys. Res. Lett.*, *32*, L19501, doi:10.1029/2005GL024057.
- Yang, D., X. Shi, and P. Marsh (2014), Variability and extreme of Mackenzie River daily discharge during 1973–2011, *Quat. Int.*, *380–381*, 159–168, doi:10.1016/j.quaint.2014.09.023.
- Ye, B., D. Yang, Z. Zhang, and D. L. Kane (2009), Variation of hydrological regime with permafrost coverage over Lena Basin in Siberia, *J. Geophys. Res.*, *114*, D07102, doi:10.1029/2008JD010537.
- Ye, H., D. Yang, and D. Robinson (2008), Winter rain on snow and its association with air temperature in northern Eurasia, *Hydrol. Processes*, *22*(15), 2728–2736.
- Zhang, H., G. H. Huang, D. Wang, and X. Zhang (2011), Uncertainty assessment of climate change impacts on the hydrology of small prairie wetlands, *J. Hydrol.*, *396*(1–2), 94–103, doi:10.1016/j.jhydrol.2010.10.037.
- Zhang, K., J. S. Kimball, Q. Mu, L. A. Jones, S. J. Goetz, and S. W. Running (2009), Satellite based analysis of northern ET trends and associated changes in the regional water balance from 1983 to 2005, *J. Hydrol.*, *379*(1), 92–110.
- Zhang, X., A. Sorteberg, J. Zhang, R. Gerdes, and J. C. Comiso (2008), Recent radical shifts of atmospheric circulations and rapid changes in Arctic climate system, *Geophys. Res. Lett.*, *35*, L22701, doi:10.1029/2008GL035607.
- Zhang, X., J. He, J. Zhang, I. Polyakov, R. Gerdes, J. Inoue, and P. Wu (2013), Enhanced poleward moisture transport and amplified northern high-latitude wetting trend, *Nat. Clim. Change*, *3*(1), 47–51, doi:10.1038/nclimate1631.